arXiv:1210.6117v1 [hep-ph] 23 Oct 2012

Calculation of binding energies and masses of quarkonia in analytic QCD models

César Ayala[1]* and Gorazd Cvetič[12]†
[1]*Department of Physics, Universidad Técnica Federico Santa María (UTFSM), Valparaíso, Chile*
[2]*Centro Científico-Tecnológico de Valparaíso, UTFSM, Chile*

(Dated: October 24, 2012)

We extract quark masses $\overline{m}_q$ $(q = b, c)$ from the evaluation of the masses of quarkonia $\Upsilon(1S)$ and $J/\psi(1S)$, performed in two analytic QCD models, and in perturbative QCD in two renormalization schemes. In analytic QCD the running coupling has no unphysical singularities in the low-momentum regime. We apply the Minimal Analytic (MA) model of Shirkov, Solovtsov, et al.; and the 2-delta analytic model (2danQCD). The latter, in contrast to MA, at higher energies basically coincides with the perturbative QCD (in the same scheme). We use the renormalon-free mass $\overline{m}_q$ as input. The separation of the soft and ultrasoft parts of the binding energy $E_{q\bar{q}}$ is performed by the requirement of the cancellation of the leading infrared renormalon. The analysis in 2danQCD indicates that the low-momentum ultrasoft regime is important for the extraction of the masses $\overline{m}_q$, especially $\overline{m}_b$. 2danQCD gives us clues on how to estimate the influence of the ultrasoft sector on $\overline{m}_q$ in general. These effects lead to relatively large values $\overline{m}_b \approx 4.35$ GeV in 2danQCD. In perturbative QCD in $\overline{\rm MS}$ scheme these effects are even stronger and give larger uncertainties in $\overline{m}_b$. The MA model gives small ultrasoft effects and the extracted values of $\overline{m}_b$ agree with those in most of the literature ($\overline{m}_b \approx 4.2$ GeV). The extracted values of $\overline{m}_c$ in all four models are about 1.26-1.27 GeV and agree well with those in the literature.

PACS numbers: 12.38.Cy, 12.38.Aw,12.40.Vv

I. INTRODUCTION

Most of the calculations of the masses of heavy quarkonia are based on perturbative expansions. These expansions come from the knowledge of the static quark-antiquark potential $V(r)$ which has been calculated up to N^{2}LO ($\sim \alpha_s^3/r$) in Refs. [1, 2]. At N^{3}LO ($\sim \alpha_s^4/r$) level, ultrasoft gluons contribute [3, 4] to $V(r)$ and the calculation of these terms has been completed in Refs. [5, 6]. The expansion coefficients for the mass of the (heavy) $q\bar{q}$ quarkonium vector $(S = 1)$ or scalar $(S = 0)$ ground state (i.e., with: $n = 1$ and $\ell = 0$) are given, for example, in Ref. [7], for the terms including $\sim m_q \alpha_s^5$. The latter term is completely known because the parameter a_3 in the corresponding aforementioned $\sim \alpha_s^4$ contribution to the static potential is now known, Refs. [5, 6]. For a review of the topic, see, e.g. Ref. [8].

Most of the radiative contributions to the quarkonium mass expansion are from the so called soft sector of gluon momenta, $Q \sim m_q \alpha_s$.[1] In the case of $b\bar{b}$ and $c\bar{c}$, this is around 2 GeV and 1 GeV, respectively. At $Q \approx 2$ GeV scales, the perturbative QCD (pQCD) coupling $\alpha_s(Q^2)$ is marginally reliable; at $Q \approx 1$ GeV it is unreliable. This is so because the pQCD coupling $\alpha_s(Q^2)$ suffers from unphysical (Landau) singularities at low spacelike q^2 $(\equiv -Q^2)$, i.e., at $0 < Q^2 < \Lambda^2$ where $\Lambda^2 \sim 10^{-1}$ GeV^2; for $Q^2 \sim 1$ GeV^2 it is dangerously close to these singularities, thus unreliable.

The aforementioned Landau singularities are not physical because they do not possess the analytic properties of the spacelike observables $\mathcal{D}(Q^2)$ (such as Adler function), the latter properties being dictated by the general properties of quantum field theories [9, 10] including causality. Namely, $\mathcal{D}(Q^2)$ must be an analytic function in the entire Q^2-complex plane, with the exception of the timelike semiaxis $Q^2 < -M_{\rm thr}^2$, where $M_{\rm thr} \sim 10^{-1}$ GeV is a particle production threshold. Specifically, the evaluation of spacelike quantities $\mathcal{D}(Q^2)$ in pQCD, as a (truncated) power series of the perturbative running coupling (couplant) $a_{\rm pt}(\kappa Q^2) \equiv \alpha_s(\kappa Q^2)/\pi$ (with $\kappa \sim 1$), does not respect these important analytic properties of $\mathcal{D}(Q^2)$. These problems, within the context of QCD, were first addressed in the seminal works of Shirkov, Solovtsov *et al.* [11–14]. There, the perturbative running coupling $a_{\rm pt}(Q^2) \equiv \alpha_s(Q^2)/\pi$ was made an analytic (in the aforementioned sense) function of Q^2, $a_{\rm pt}(Q^2) \mapsto \mathcal{A}_1^{\rm (MA)}(Q^2)$, by keeping in the dispersion relation the discontinuity function $\rho_1^{\rm (pt)}(\sigma) \equiv {\rm Im}\ a_{\rm pt}(Q^2 = -\sigma - i\epsilon)$ unchanged on the entire negative axis in the

---

*Electronic address: c.ayala86@gmail.com
†Electronic address: gorazd.cvetic@usm.cl

[1] The momentum transfer q of the gluon is spacelike, i.e., $q^2 \equiv -Q^2 < 0$, because the scattering of the quark and antiquark in the quarkonium is of the t-channel type.

complex Q^2-plane (i.e., for $\sigma \geq 0$), and was set equal to zero on the nonphysical cut $0 < Q^2 < \Lambda^2$. For this reason, this analytic QCD (anQCD) model will be referred to here as Minimal Analytic (MA).[2]

Various other analytic QCD (anQCD) models for $\mathcal{A}_1(Q^2)$ can be constructed, and have been presented in the literature, among them Refs. [15–23]. These models fulfill certain additional constraints at low and/or at high Q^2. For further literature on various analytic QCD models, we refer to review articles Refs. [24–27]. Some newer constructions of anQCD models of $\mathcal{A}_1(Q^2)$ include those based on specific classes of β functions with nonperturbative contributions [28] or without such contributions [29, 30], and those based on modifications of the discontinuity function $\rho_1^{(\rm pt)} \mapsto \rho_1$ [$\rho_1(\sigma) \equiv {\rm Im}\ \mathcal{A}_1(Q^2 = -\sigma - i\epsilon)$] at low (positive) σ where ρ_1 is parametrized in a specific manner (with delta functions), cf. Refs. [31, 32].

The model of Ref. [32], an extension of the model of Ref. [31], will be called 2-delta analytic QCD (2danQCD), because its discontinuity function $\rho_1(\sigma)$ is parametrized in the unknown low-σ regime by two delta functions. It has a specific attractive feature that in the high-scale regime ($|Q^2| > \Lambda^2$) it practically coincides with the corresponding (in the scheme) pQCD coupling: $\mathcal{A}_1(Q^2) - a_{\rm pt}(Q^2) \sim (\Lambda^2/Q^2)^5$. This makes it possible for the model to be applied with Operator Product Expansion (OPE) [33] including terms of dimension $D \leq 8$ without the need to modify the ITEP School interpretation of the OPE [34, 35]. The latter basically states that the terms in OPE of higher dimension ($D > 0$) originate from the infrared regime only.

Once having an anQCD model, i.e., a model for the analytic analog $\mathcal{A}_1(Q^2)$ of the pQCD coupling $a_{\rm pt}(Q^2)$ (or, equivalently, for the discontinuity function $\rho_1(\sigma)$), the analytic analogs $\mathcal{A}_n(Q^2)$ of the higher powers $a_{\rm pt}(Q^2)^n$ have to be constructed from it. In the case of MA/APT ($\mathcal{A}_1^{(\rm MA)}(Q^2)$), the couplings $\mathcal{A}_n^{(\rm MA)}(Q^2)$ were constructed in Ref. [36] for integer n and in Refs. [37–39] for noninteger n (cf. also Ref. [27]). In the general case of analytic $\mathcal{A}_1(Q^2)$, the quantities $\mathcal{A}_n(Q^2)$ were constructed in Refs. [20, 21] for integer n, as linear combinations of logarithmic derivatives $\widetilde{\mathcal{A}}_k(Q^2) \propto d^{k-1}\mathcal{A}_1(Q^2)/d(\ln Q^2)^{k-1}$ ($k \geq n$),[3] and were extended to noninteger n in Ref. [41].

In this work, we apply two anQCD models, MA/APT of Refs. [11–14], and 2danQCD of Refs. [32, 33], to evaluations of the perturbation series of the binding energy $E_{q\bar{q}}$ of heavy quarkonia ($\Upsilon(1S)$ and $J/\psi(1S)$) and of the quark pole mass m_q, and thus to evaluations of the masses of these quarkonia $M_{q\bar{q}} = 2m_q + E_{q\bar{q}}$. As input parameter we use the renormalon-free quark mass $\overline{m}_q$ ($\equiv \overline{m}_q(\mu^2 = \overline{m}_q^2)$) of the corresponding quark $q = b, c$ (also called $\overline{\rm MS}$ quark mass), and the anQCD coupling of the model. Since the quarkonia masses are well measured, we can extract the values of $\overline{m}_q$. We also perform the same analysis in pQCD in the corresponding renormalization schemes.

In Sec. II we briefly describe the MA/APT model (Subsec. II A) and the 2danQCD model (Subsec. II B). In Sec. III we present the procedures of evaluation of the binding energy $E_{q\bar{q}}$ and of the quark pole mass m_q, in terms of the mass $\overline{m}_q$ and of the couplings. Furthermore, we explain how the cancellation of the leading infrared renormalon in the sum $2m_q + E_{q\bar{q}}$ allows us to separate the ultrasoft from the soft part of the binding energy. The numerical results and the extractions of the masses $\overline{m}_b$ and $\overline{m}_c$ are presented in Sec. IV (Subsections IV A and IV B, respectively). The evaluations are performed in the two aforementioned analytic models MA/APT and 2danQCD, and in pQCD in the corresponding two renormalization schemes ($\overline{\rm MS}$, and in the scheme of 2danQCD called here the Lambert scheme). In Sec. V we summarize our results and draw certain conclusions. Appendix A contains the expressions of the coefficients of the perturbation expansion of $E_{q\bar{q}}$, available from the literature. Appendix B contains a renormalon-based estimation of the coefficient at $a_{\rm pt}^4$ term in the perturbation expansion of $m_q/\overline{m}_q$.

II. TWO ANALYTIC QCD MODELS

A. Minimal Analytic (MA/APT) model

As already mentioned in the Introduction, the coupling $a_{\rm pt}(Q^2) \equiv \alpha_s(Q^2)/\pi$ in pQCD in the usual renormalization schemes (such as $\overline{\rm MS}$, or the $\beta_2 = \beta_3 = \ldots = 0$ scheme), possesses unphysical (Landau) singularities inside the required analyticity regime $Q^2 \in \mathbb{C}\backslash(-\infty, 0]$. Specifically, it has a cut on the semiaxis $(-\infty, \Lambda_{\rm L}^2)$ (where $\Lambda_{\rm L}^2 \sim 10^{-1}$ GeV^2 is the "Landau" branching point), thus offending the analyticity requirement on the cut sector $Q^2 \in (0, \Lambda_{\rm L}^2)$. Application of the Cauchy theorem gives us for the power $a_{\rm pt}^{\nu}(Q^2)$ ($\nu > 0$) of this spacelike coupling the following dispersion relation:

---

[2] In the literature, it is also referred to as Analytic Perturbation Theory (APT). We refer to it as MA, or MA/APT.

[3] The relations between such functions $\widetilde{\mathcal{A}}_k$'s and $\mathcal{A}_n$'s, allowing a recurrent construction of $\mathcal{A}_n$, for integer n within the context of the MA/APT model, were given also in Refs. [25, 40].

$$a_{\rm pt}^{\nu}(Q^2) = \frac{1}{\pi} \int_{\sigma=-\Lambda_{\rm L}^2-\eta}^{\infty} \frac{d\sigma \ {\rm Im} \ a_{\rm pt}^{\nu}(-\sigma - i\epsilon)}{(\sigma + Q^2)}, \tag{1}$$

and $\eta \to +0$. Eliminating the aforementioned cut sector, and keeping the rest of the discontinuity function $\rho_1^{\rm (pt)}(\sigma) \equiv {\rm Im} \ a_{\rm pt}(Q^2 = -\sigma - i\epsilon)$ unchanged, results in the MA/APT model [11–14], i.e., the spacelike coupling

$$\left(a_{\rm pt}^{\nu}(Q^2) \right)_{\rm an}^{\rm (MA)} \equiv \mathcal{A}_{\nu}^{\rm (MA)}(Q^2) = \frac{1}{\pi} \int_{\sigma=0}^{\infty} d\sigma \frac{{\rm Im} \ a_{\rm pt}^{\nu}(-\sigma - i\epsilon)}{(\sigma + Q^2)} \ . \tag{2}$$

In evaluation of general spacelike observables in pQCD, terms of the type $a_{\rm pt}^{\nu}(Q^2) \ln^k a_{\rm pt}(Q^2) \ln^{\ell}(Q^2/m^2)$ may appear, with $\nu > 0$ and (nonnegative) integers k and ℓ. In MA/APT, in principle, the factor $\ln^{\ell}(Q^2/m^2)$ may be included in the MA-type analytization (see, for example, Ref. [27]). However, the factor $\ln^{\ell}(Q^2/m^2)$ is analytic function in the Q^2-complex plane with the exclusion of the nonpositive axis $Q^2 \leq 0$. On the other hand, this is also the regime of analyticity of $\mathcal{A}_1^{\rm (MA)}(Q^2)$ [note that $Q^2 = 0$ is a point of nonanalyticity of $\mathcal{A}_1^{\rm (MA)}(Q^2)$ because $d\mathcal{A}_1^{\rm (MA)}(Q^2)/dQ^2 = \infty$ there]. Therefore, we will not include the factor $\ln^{\ell}(Q^2/m^2)$ in the (MA-type) of analytization. With these considerations, all the other pQCD terms in such observables, $a_{\rm pt}^{\nu}(Q^2) \ln^k a_{\rm pt}(Q^2)$, are made analytic in MA/APT by the same procedure as $a_{\rm pt}^{\nu}(Q^2)$ in Eq. (2)

$$\begin{aligned} \left(a_{\rm pt}^{\nu}(Q^2) \ln^k a_{\rm pt}(Q^2) \right)_{\rm an}^{\rm (MA)} &\equiv \mathcal{A}_{\nu,k}^{\rm (MA)}(Q^2) \\ = \frac{1}{\pi} \int_{\sigma=0}^{\infty} d\sigma & \frac{{\rm Im} \left[a_{\rm pt}^{\nu}(-\sigma - i\epsilon) \ln^k a_{\rm pt}(-\sigma - i\epsilon) \right]}{(\sigma + Q^2)} \ . \end{aligned} \tag{3}$$

In this context, the authors of Refs. [27, 37–39] derived explicit simple formulas for these MA expressions at one-loop level, and the related more involved expressions at higher-loop levels – expressions from which the analyticity structure in Q^2 and in ν can be more clearly seen. For our calculational purposes, though, the dispersion formulas (3) will be applied, numerically, when the MA/APT model is used.

In general analytic QCD (anQCD) models, for any spacelike observable $\mathcal{D}(Q^2)$ which in pQCD is expanded in powers of $a_{\rm pt}^{\nu}(\mu^2) \ln^k a_{\rm pt}(\mu^2)$ (where $\mu^2 \sim Q^2$ is the renormalization scale), the analytization method consists in principle in a replacement

$$a_{\rm pt}^{\nu}(\mu^2) \ln^k a_{\rm pt}(\mu^2) \mapsto \left(a_{\rm pt}^{\nu}(\mu^2) \ln^k a_{\rm pt}(\mu^2) \right)_{\rm an} \equiv \mathcal{A}_{\nu,k}(\mu^2) \ . \tag{4}$$

The explicit procedure of such analytization, in a general anQCD model, is described in Ref. [41].

MA/APT has a relatively specific and interesting property, namely, that the couplings differ from the corresponding pQCD couplings in a nonnegligible way even at high $|Q^2|$

$$\mathcal{A}_1^{\rm (MA)}(Q^2) - a_{\rm pt}(Q^2) \sim \Lambda^2/Q^2 \qquad (|Q^2| > \Lambda^2) \ . \tag{5}$$

As a consequence, $\mathcal{A}_n^{\rm (MA)}(Q^2)$ and $a_{\rm pt}(Q^2)^n$ differ from each other in a nonnegligible way even at high $|Q^2|$ [14]; we cannot approximate $\mathcal{A}_1^{\rm (MA)}(Q^2)$ well with $a_{\rm pt}(Q^2)$ even at as high $|Q^2|$ as $\sim M_Z^2$. The (only) free parameter in MA/APT is the $\overline{\rm MS}$ scale $\overline{\Lambda}$, which appears in the usual perturbation expansion of the underlying pQCD coupling $a_{\rm pt}(Q^2; \overline{\rm MS})$ in inverse powers of $\ln(Q^2/\overline{\Lambda}^2)$ [i.e., the $a_{\rm pt}$ whose discontinuity functions appear in the relations (2)-(3)].

It turns out that for MA/APT this scale varies in the interval $\overline{\Lambda}_{N_f=5} = 0.260 \pm 0.030$ GeV. The analysis of high energy QCD observables (with $|Q| \gtrsim 10^1$ GeV) in Ref. [14] suggests a value of $\overline{\Lambda}_{N_f=5} \approx 0.290$ GeV [corresponding to $\alpha_s(M_Z^2, \overline{\rm MS}) \approx 0.124$]. The works of Refs. [27, 38, 39] use the value $\overline{\Lambda}_{N_f=5} \approx 0.260$ GeV instead [corresponding to $\alpha_s(M_Z^2, \overline{\rm MS}) \approx 0.122$ and giving the timelike (Minkowskian) coupling $\mathfrak{A}_2(M_Z^2) \approx 0.120$]. On the other hand, if the "average" e^+e^- annihilation ratio $R(s)$ value at $\sqrt{s} = M_Z$ is taken to be ≈ 1.03904 as used, e.g., in Ref. [42], MA/APT approach of evaluating $R(M_Z^2)$ gives $\overline{\Lambda}_{N_f=5} \approx 0.225$ GeV [43].

In comparison, pQCD analyses give for $\alpha_s(M_Z^2)$ the world average 0.1184 ± 0.0007 [44], corresponding to $\overline{\Lambda}_{N_f=5} = 0.213 \pm 0.008$ GeV, i.e., significantly lower scales than for MA/APT.[4]

---

[4] Also the anQCD models of Refs. [15, 16] differ nonnegligibly from pQCD, even at high $|Q^2|$.

Due to the nonnegligible difference, Eq. (5), the threshold effects of heavy quarks have to be implemented in MA/APT. This is performed by requiring the continuity of the perturbative coupling [whose discontinuities are used in MA/APT]

$$a_{\rm pt}(Q^2) \equiv f_{N_f}(Q^2/\overline{\Lambda}^2_{N_f}) \ , \tag{6}$$

at positive "threshold" values $Q^2 = Q^2_{N_f - 1 \mapsto N_f} > 0$ chosen to be for $N_f = 4, 5, 6$ the squares of the current heavy quark masses $m_c \equiv m_4$, $m_b \equiv m_5$ and $m_t \equiv m_6$, respectively [14]

$$f_{N-1}(m_N^2/\overline{\Lambda}^2_{N-1}) = f_N(m_N^2/\overline{\Lambda}^2_N) \qquad (N=4,5,6) \ . \tag{7}$$

Further, it is assumed that, for complex Q^2, the values of $a_{\rm pt}(Q^2)$ involve the scale $\overline{\Lambda}_{N_f}$ determined by the absolute value $|Q^2|$

$$a_{\rm pt}(Q^2) = f_N(Q^2/\overline{\Lambda}^2_N) \qquad \text{for} : m_N^2 < |Q^2| < m_{N+1}^2 \ . \tag{8}$$

Thus constructed $a_{\rm pt}(Q^2)$ gives the piecewise dicontinuous global MA/APT discontinuity functions. For example, for $\rho^{(\rm pt)}_{\nu,k}(\sigma) \equiv {\rm Im}[a_{\rm pt}^{\nu} \ln^k a_{\rm pt}(-\sigma - i\epsilon)]$, this globalization results in

$$\begin{aligned} \rho^{(\rm pt)}_{\nu,k}(\sigma) &\equiv {\rm Im} \left(a_{\rm pt}^{\nu}(-\sigma - i\epsilon) \ln^k a_{\rm pt}(-\sigma - i\epsilon) \right) \\ &= {\rm Im} \left[f_N(z)^{\nu} \ln^k f_N(z) \right] \Big|_{z = -\sigma/\overline{\Lambda}^2_N - i\epsilon} \qquad \text{for} : (m_N^2 < \sigma < m_{N+1}^2) \ . \end{aligned} \tag{9}$$

Despite the discontinuity of $\rho^{(\rm pt)}_{\nu,k}(\sigma)$, the MA/APT analytic analogs (3) are analytic functions.

For the underlying $a_{\rm pt}(Q^2) \equiv \alpha_s(Q^2)/\pi$ we will use the effective three-loop coupling of the form [45–47]

$$a_{\rm pt}(Q^2) = -\frac{1}{c_1} \frac{1}{\left[1 - c_2/c_1^2 + W_{\mp 1}(z)\right]} \ , \tag{10}$$

where $c_j = \beta_j/\beta_0$ $(j=1,2)$, $Q^2 = |Q^2| \exp(i\phi)$, and W_{-1} and W_{+1} are the branches of the Lambert function for the case $0 \le \phi < +\pi$ and $-\pi < \phi < 0$, respectively, and

$$z(Q^2) = -\frac{1}{c_1 e} \left(\frac{|Q^2|}{\Lambda^2} \right)^{-\beta_0/c_1} \exp\left(-i \beta_0 \phi/c_1\right) \ . \tag{11}$$

We call the scale Λ^2 (~ 0.1 GeV^2) appearing here the Lambert scale. The aforementioned values $\overline{\Lambda}_{N_f=5} = 0.260 \pm 0.030$ GeV correspond to $\Lambda_{N_f=5} = 0.438 \pm 0.051$ GeV. The solution (10) is a solution to the renormalization group equation (RGE) where the beta function has the Padé form $\beta(a_{\rm pt}) \propto a_{\rm pt}^2 [1/1](a_{\rm pt})$

$$\frac{d a_{\rm pt}(Q^2)}{d \ln Q^2} = -\beta_0 a_{\rm pt}^2 \frac{\left[1 + (c_1 - (c_2/c_1)) a_{\rm pt}\right]}{\left[1 - (c_2/c_1) a_{\rm pt}\right]} \ . \tag{12}$$

The expansion of beta function $\beta(a_{\rm pt})$ on the right-hand side gives

$$\beta(a_{\rm pt}) = -\beta_0 a_{\rm pt}^2 \left(1 + c_1 a_{\rm pt} + c_2 a_{\rm pt}^2 + c_3 a_{\rm pt}^3 + \ldots \right) \ , \tag{13}$$

where the higher renormalization scheme parameters c_j $(j \ge 3)$ are fixed by the leading scheme parameter c_2: $c_j = c_2^{j-1}/c_1^{j-2}$. The three-loop scheme parameter c_2 will be fixed so that it is equal to the $\overline{\rm MS}$ value $\overline{c}_2$ (for a given N_f). On the other hand, $\beta_0 = (1/4)(11 - 2N_f/3)$ and $c_1 = \beta_1/\beta_0 = (1/4)(102 - 38N_f/3)/(11 - 2N_f/3)$ are universal constants. The four-loop scheme parameter $c_3 \equiv \beta_3/\beta_0$ is equal here to c_2/c_1^2, as can be seen from Eq. (13), i.e., the value of c_3 cannot be adjusted (e.g., to the $\overline{\rm MS}$ value $\overline{c}_3$) once the value of c_2 has been fixed. The specific choice (10) is convenient because it represents an explicit function[5] and it is thus easy to evaluate it and the corresponding discontinuity functions $\rho^{(\mathrm{pt})}_{\nu,k}(\sigma)$, Eq. (3), in practice. On the other hand, it approximates the usual ($\overline{\mathrm{MS}}$) $a_{\mathrm{pt}}(Q^2)$ to three-loops. It has Landau singularities (this is true even for any values of c_2).

---

[5] In Mathematica [48] the Lambert function $W_n(z)$ is implemented under the name ProductLog$[n, z]$.

Using the aforementioned value $\Lambda_5 = 0.438 \pm 0.051$ GeV ($\overline{\Lambda}_5 = 0.260 \pm 0.030$ GeV), the continuity conditions (7) give us at other values of N_f: $\Lambda_4 = 0.745 \pm 0.077$ GeV ($\overline{\Lambda}_4 = 0.351 \pm 0.036$ GeV), $\Lambda_3 = 0.948 \pm 0.085$ GeV ($\overline{\Lambda}_3 = 0.382 \pm 0.033$ GeV), and $\Lambda_6 = 0.119 \pm 0.015$ GeV ($\overline{\Lambda}_6 = 0.112 \pm 0.014$ GeV). The first three quark flavors are regarded to be massless.

In this context, we mention that the elimination of the Landau singularities (i.e., analytization) of $a_{\mathrm{pt}}(Q^2)$ can be performed in various ways, not just in the "minimal" way of Eq. (2). This analytization, in general results in an (analytic) running coupling $\mathcal{A}_1(Q^2)$ which differs from the corresponding perturbative one by a power term

$$\mathcal{A}_1(Q^2) = a_{\mathrm{pt}}(Q^2) + \mathcal{O}\left(\left(\frac{\Lambda^2}{Q^2}\right)^n\right) \ , \quad (Q^2 > \Lambda^2) \ . \tag{14}$$

In the case of MA/APT, the power index is $n = 1$, cf. Eq. (5). This index can be increased to $n = 3$ [17, 31], and even to $n = 5$ [32], while still maintaining analyticity. In Refs. [29, 30] the problem of finding perturbative analytic couplings (in specific schemes) was investigated. It turned out that such couplings always gave significantly too low value of the (strangeless) semihadronic tau lepton decay ratio r_τ, unless the scheme was changed in a drastic way which made the perturbation series diverge strongly after the first four terms.

Therefore, the appearance of the power terms as given in Eq. (14) seems to be a general feature of procedures which eliminate the unphysical (Landau) singularities of the coupling in the complex Q^2-plane. These power terms are of ultraviolet origin and thus contravene the philosophy of the ITEP school [34], according to which the power terms in (inclusive) QCD observables appear in the Operator Product Expansion (OPE) and are of infrared origin. Furthermore, in MA/APT we have $n = 1$ and the terms $\sim \Lambda^2/Q^2$ appear even in the massless QCD, i.e., in the case when such terms are not allowed in the usual pQCD+OPE approach for observables related with the vacuum expectation values (such as the Adler function). In general, the ITEP interpretation of OPE must be abandoned in those anQCD models in which the index n in Eq. (14) is low (e.g., $n \leq 2$), and modified/restricted in others (e.g., when $n = 3, 4$).[6]

Despite these reservations, we evaluate in this work quarkonium masses with the MA/APT model of anQCD, in order to illustrate the effects of elimination of Landau singularities, within such a context, on the behavior of the series. MA/APT model is also relatively simple to apply technically.

We will also apply another anQCD model to the evaluation of quarkonium masses, namely the so called two-delta anQCD [32], in which the index in Eq. (14) is $n = 5$. Evaluations in such models are technically more demanding, though, because the evaluation of the analytic analogs of noninteger powers a_{pt}^{ν} and of the logarithm-type terms $a^n \ln^k a$ is more involved [41].

B. Two-delta analytic QCD model

It is possible to construct such anQCD models which, at high momenta, practically merge with pQCD; namely, with the analytic coupling $\mathcal{A}_1(Q^2)$ which fulfills Eq. (14) with $n > 1$. One such model, with $n = 3$, was constructed in Ref. [17][7] Another such anQCD model, which we can call one-delta model, was constructed in Ref. [31], and also satisfies Eq. (14) with $n = 3$. The idea was simple: the perturbative discontinuity function $\rho_1^{(\mathrm{pt})}(\sigma) \equiv \mathrm{Im}\, a_{\mathrm{pt}}(Q^2 = -\sigma - i\epsilon)$, for $\sigma > 0$, was kept unchanged for σ down to a "pQCD-onset-scale" $M_0^2 \sim 1$ GeV^2, and the unknown low-energy regime $0 < \sigma < M_0^2$ was parametrized with one delta function at a scale M_1^2 (such that: $0 < M_1^2 < M_0^2$)[8]

$$\rho_1^{(\mathrm{1d})}(\sigma) = \pi f_1^2 \Lambda^2 \delta(\sigma - M_1^2) + \Theta(\sigma - M_0^2) \ , \rho_1^{(\mathrm{pt})}(\sigma) \ , \tag{15}$$

where for $a_{\mathrm{pt}}(Q^2)$ [the latter defining $\rho_1^{(\mathrm{pt})}(\sigma)$] the renormalization scheme with $c_2 = c_3 = \ldots = 0$ was used. The (Lambert) scale parameter Λ was fixed by the value of $\alpha_s(M_Z^2, \overline{\mathrm{MS}})$. The dimensionless parameters $s_j \equiv M_j^2/\Lambda^2$

---

[6] Nonetheless, even in MA/APT, OPE can be applied, and has successfully been applied – see, for example, the inclusion of higher-twist terms in the MA/APT vs pQCD analysis of the Bjorken polarized sum rule at low Q^2, Refs. [49]. We note that the leading-twist ($D = 0$) term, in such a MA/APT+OPE approach, contains implicitly (parts of) the power term contributions $\sim (\Lambda^2/Q^2)^n$, $n = 1, 2, \ldots$.

[7] The higher couplings $\mathcal{A}_n(Q^2) = (a_{\mathrm{pt}}(Q^2)^n)_{\mathrm{an}}$ were not constructed there, though.

[8] A similar idea was applied in Refs. [50, 51] directly to spectral functions of the vector current correlators. Other approaches of eliminating unphysical singularities directly from specific (spacelike) obsevables, were presented in Refs. [52–54].

TABLE I: Values of the parameters of the considered 2danQCD model. We consider $c_2 = -4.76$ ($M_0 = 1.25$ GeV) as the central representative case. The Lambert scale values in the corresponding cases are for the QCD coupling parameter value $\alpha_s^{(\overline{\rm MS})}(M_Z^2) = 0.1184$.

$c_2 = \beta_2/\beta_0$	s_0	s_1	f_1^2	s_2	f_2^2	Λ [GeV]	M_0	$\mathcal{A}_1(0)$
-2.10	17.09	12.523	0.1815	0.7796	0.3462	0.363	1.50	0.544
-4.76	23.06	16.837	0.2713	0.8077	0.5409	0.260	1.25	0.776
-5.73	25.01	18.220	0.3091	0.7082	0.6312	0.231	1.15	1.00

($j = 0, 1$) and f_1^2 were then fixed by the requirement of $n = 3$ in Eq. (14) (these are in fact two conditions) and by the requirement that the central experimental value of the $(V + A)$ τ-decay ratio r_τ, namely $(r_\tau)_{\rm exp} = 0.203 \pm 0.004$ [55, 56],[9] be reproduced in the model.

As argued at the end of the previous Subsection, the ITEP interpretation of the OPE requires that index n in Eq. (14) be relatively high,[10] $4 < n < 9$. If, instead of one delta, we use two deltas to parametrize the otherwise unknown low-σ regime behavior of the spectral function $\rho_1(\sigma)$, we obtain an analytic QCD model with $\mathcal{A}_1(Q^2)$ fulfilling the relation (14) with $n = 5$, Ref. [32],

$$\rho_1^{(2{\rm d})}(\sigma; c_2) = \pi \sum_{j=1}^{2} f_j^2 \Lambda^2 \, \delta(\sigma - M_j^2) + \Theta(\sigma - M_0^2) \times \rho_1^{(\rm pt)}(\sigma; c_2) \tag{16}$$

$$= \pi \sum_{j=1}^{2} f_j^2 \, \delta(s - s_j) + \Theta(s - s_0) \times r_1^{(\rm pt)}(s; c_2) \; . \tag{17}$$

Here, the dimensionless squared energy $s = \sigma/\Lambda^2$ was introduced, and five dimensionless parameters: $s_j = M_j^2/\Lambda^2$ ($j = 0, 1, 2$) and f_k^2 ($k = 1, 2$). Further, $c_2 = \beta_2/\beta_0$ is the scheme parameter, and $r_1^{(\rm pt)}(s; c_2) = \rho_1^{(\rm pt)}(\sigma; c_2) = {\rm Im}\, a_{\rm pt}(Q^2 = -\sigma - i\epsilon; c_2)$ is the perturbative spectral function of $a_{\rm pt}(Q^2)$ in terms of the dimensionless energy variable $s = \sigma/\Lambda^2$, where $a_{\rm pt}(Q^2)$ is given by Eqs. (10)-(11) [see also Eqs. (12)-(13)]. The spacelike coupling $\mathcal{A}_1(Q^2)$ is then obtained by the dispersion relation

$$\mathcal{A}_1^{(2{\rm d})}(Q^2) = \frac{1}{\pi} \int_{\sigma=0}^{\infty} d\sigma \frac{\rho_1^{(2{\rm d})}(\sigma)}{(\sigma + Q^2)} = \sum_{j=1}^{2} \frac{f_j^2}{(u + s_j)} + \frac{1}{\pi} \int_{s_0}^{\infty} ds \, \frac{r_1^{(\rm pt)}(s; c_2)}{(s + u)} \; , \tag{18}$$

where we denoted $u = Q^2/\Lambda^2$. This two-delta analytic QCD model (2danQCD) is to be applied in the low-momentum regime, i.e., where $n_f = 3$ (considering the current masses of u, d and s zero): $|Q^2| < (2m_c)^2$ (≈ 6.45 GeV^2). At higher $|Q^2| > (2m_c)^2$ (where $n_f \geq 4$) it is replaced by the underlying pQCD of the coupling $a_{\rm pt}(Q^2)$, Eq. (10). The Lambert scale $\Lambda^2 = \Lambda^2_{n_f=3}$ (~ 0.1 GeV^2), which is the only dimensional parameter of the model, is determined by the world average value $a_{\rm pt}(M_Z^2; \overline{\rm MS}) = (0.1184 \pm 0.0007)/\pi$, [44]. For example, when $c_2 = -4.76$, we obtain $\Lambda \approx (0.260 \pm 0.008)$ GeV. On the other hand, the five aforementioned dimensionless parameters s_j and f_k^2 are fixed, independently of the value of Λ, by the condition Eq. (14) with $n = 5$ (these are four conditions) and by the condition that the aforementioned central experimental value of the $(V + A)$ τ-decay ratio r_τ be reproduced by the model, namely $r_\tau = 0.203$. More specifically, the (four) conditions Eq. (14) (with $n = 5$) determine the four relations $s_j(s_0)$ and $f_j^2(s_0)$ ($j = 1, 2$), and the value of s_0 ($\equiv M_0^2/\Lambda^2$) is then determined (if Λ is already fixed) by the condition $r_\tau(V + A) = 0.203$. The (scheme) parameter c_2 remains a free parameter. Varying c_2 results in variation of the pQCD-onset scale M_0 ($\equiv \sqrt{s_0}\Lambda$) and in variation of $\mathcal{A}_1(Q^2)$ at low $|Q^2|$. For phenomenological reasons, we wish to have M_0 below the τ-lepton mass m_τ (≈ 1.78 GeV) and to have the largest $|\mathcal{A}_1(Q^2)| < 1.0$. More specifically, we vary c_2 in such a way that $1.0 \text{ GeV} \leq M_0 \leq 1.5$ GeV and $\mathcal{A}_1(0) < 1.0$. This results in the central value $c_2 = -4.46$ for which $M_0 = 1.25$ GeV (and $\mathcal{A}_1(0) = 0.776$), and two border values: $c_1 = -2.10$ for which $M_0 = 1.50$ GeV (and $\mathcal{A}_1(0) = 0.544$); $c_2 = -5.73$ for which $\mathcal{A}_1(0) = 1.00$ (and $M_0 = 1.15$ GeV).[11] The model parameters for the central and the two border cases are given in Table I.

---

[9] r_τ is the QCD part of the V+A decay ratio $R_\tau(\Delta S = 0)$, with the (small) quark mass effects subtracted. It is normalized in the canonical way: $r_\tau = a + \mathcal{O}(a^2)$.

[10] Small-size instanton effects may lead to the conditions Eq. (14) to be valid only for up to index n such that $2n$ is the largest dimension of condensates not affected by the instantons. Instanton-antiinstanton gas scenarios lead to $n < 4\beta_0$ ($= 9$ for $n_f = 3$), cf. Ref. [35].

[11] In Ref. [32] we included the case $c_2 = -7.15$, for which $M_0 = 1.0$ GeV. However, in that case, $\mathcal{A}_1(0) = 2.29$, and this indicates that the model has infrared instabilities at such low c_2 values.

Variation of the scheme parameter c_2, at fixed central world average value $a_{\rm pt}(M_Z^2;\overline{\rm MS}) = 0.1184/\pi$, implied the corresponding variation of the parameters s_j, f_k^2 and Λ in Table I, such that the relation (14) with $n=5$ is fulfilled and τ-decay ratio value $r_\tau = 0.203$ is reproduced.

On the other hand, if we vary the QCD coupling parameter within the world average interval, $a_{\rm pt}(M_Z^2;\overline{\rm MS}) = (0.1184 \pm 0.0007)/\pi$, [44], while keeping c_2 $(=-4.76)$ and all the dimensionless parameters fixed, then only the scale parameter Λ varies, $\Lambda = 0.260 \pm 0.008$ GeV, reflecting the changing $a_{\rm pt}(M_Z^2;\overline{\rm MS})$. The fulfillment of the relation (14) with $n=5$, at given c_2, fixes the values of the four dimensionless parameters s_k and f_k^2 $(k=1,2)$ in terms of the parameter s_0 $(=23.06)$. The resulting value of τ-decay ratio then varies with $a_{\rm pt}(M_Z^2;\overline{\rm MS})$: $r_\tau = 0.203 \pm 0.006$. This is acceptably compatible with the measured values $(r_\tau)_{\rm exp} = 0.203 \pm 0.004$ [55, 56].

For further details on 2danQCD model we refer to Ref. [32].

The higher order couplings $A_\nu(Q^2)$ (analogs of $a_{\rm pt}^\nu(Q^2)$) in this model are then obtained according to the construction for general anQCD models, explained in Refs. [20, 21] for integer ν, and Ref. [41] for general (noninteger) ν. Below we will summarize the basic aspects of such construction.

Since the general anQCD models, with the exception of MA/APT, have at low σ ($\lesssim 1\ {\rm GeV}^2$) different discontinuity function than the pQCD coupling $a_{\rm pt}(Q^2)$, we cannot use the MA/APT method, Eq. (3), for the construction of the analytic analogs of $a_{\rm pt}^\nu \ln^k a_{\rm pt}$. The analogs of the integer powers $a_{\rm pt}^n$ in such general models were constructed in Refs. [20, 21], where it was shown that it is imperative to construct first the analogs of the logarithmic derivatives of $a_{\rm pt}$ in the following way:[12]

$$\left(\frac{\partial^k a_{\rm pt}(Q^2)}{\partial (\ln Q^2)^k} \right)_{\rm an} = \frac{\partial^k \mathcal{A}_1(Q^2)}{\partial (\ln Q^2)^k} \quad (k=0,1,2,\ldots) \ . \tag{19}$$

In pQCD, the logarithmic derivatives

$$\widetilde{a}_{{\rm pt},k+1}(Q^2) \equiv \frac{(-1)^k}{\beta_0^k k!} \frac{\partial^k a_{\rm pt}(Q^2)}{\partial (\ln Q^2)^k} \ , \qquad (k=0,1,2,\ldots) \ , \tag{20}$$

are related with the powers of $a_{\rm pt} \equiv \alpha_s/\pi$ in the following way [using renormalization group equations (RGE) in pQCD]:

$$\widetilde{a}_{{\rm pt},2} = a_{\rm pt}^2 + c_1 a_{\rm pt}^3 + c_2 a_{\rm pt}^4 + c_3 a_{\rm pt}^5 + \cdots \ , \tag{21}$$

$$\widetilde{a}_{{\rm pt},3} = a_{\rm pt}^3 + \frac{5}{2} c_1 a_{\rm pt}^4 + \left(3 c_2 + \frac{3}{2} c_1^2 \right) a_{\rm pt}^5 + \cdots \ , \tag{22}$$

$$\widetilde{a}_{{\rm pt},4} = a_{\rm pt}^4 + \frac{13}{3} c_1 a_{\rm pt}^5 + \cdots \ , \qquad \widetilde{a}_{{\rm pt},5} = a_{\rm pt}^5 + \cdots \ , \qquad \text{etc.} \tag{23}$$

This means that the powers of $a_{\rm pt}$ are linear combinations of logarithmic derivatives

$$a_{\rm pt}^2 = \widetilde{a}_{{\rm pt},2} - c_1 \widetilde{a}_{{\rm pt},3} + \left(\frac{5}{2} c_1^2 - c_2 \right) \widetilde{a}_{{\rm pt},4} + \left(-\frac{28}{3} c_1^3 + \frac{22}{3} c_1 c_2 - c_3 \right) \widetilde{a}_{{\rm pt},5} + \cdots \ , \tag{24}$$

$$a_{\rm pt}^3 = \widetilde{a}_{{\rm pt},3} - \frac{5}{2} c_1 \widetilde{a}_{{\rm pt},4} + \left(\frac{28}{3} c_1^2 - 3 c_2 \right) \widetilde{a}_{{\rm pt},5} + \cdots \ , \tag{25}$$

$$a_{\rm pt}^4 = \widetilde{a}_{{\rm pt},4} - \frac{13}{3} c_1 \widetilde{a}_{{\rm pt},5} + \cdots \ , \qquad a_{\rm pt}^5 = \widetilde{a}_{{\rm pt},5} + \cdots \ , \qquad \text{etc.} \tag{26}$$

These relations, in conjunction with the analytization Eq. (19), imply that the analytic analogs $\mathcal{A}_k$ of powers $a_{\rm pt}^k$, in general anQCD models, can be expressed as linear combinations of the logarithmic derivatives

$$\widetilde{\mathcal{A}}_{k+1}(\mu^2) = \frac{(-1)^k}{\beta_0^k k!} \frac{\partial^k \mathcal{A}_1(\mu^2)}{\partial (\ln \mu^2)^k} \ , \qquad (k=1,2,\ldots) \ . \tag{27}$$

---

[12] If the analytization is performed in any other way, the renormalization scale and scheme dependence of the resulting truncated analytic series of any observable $\mathcal{D}(Q^2)$ will in general increase (instead of decrease) when the number of terms in the series increases, cf. [20, 21].

in the following form:

$$\mathcal{A}_2 = \widetilde{\mathcal{A}}_2 - c_1\widetilde{\mathcal{A}}_3 + \left(\frac{5}{2}c_1^2 - c_2\right)\widetilde{\mathcal{A}}_4 + \left(-\frac{28}{3}c_1^3 + \frac{22}{3}c_1c_2 - c_3\right)\widetilde{\mathcal{A}}_5 + \cdots \ , \tag{28}$$

$$\mathcal{A}_3 = \widetilde{\mathcal{A}}_3 - \frac{5}{2}c_1\widetilde{\mathcal{A}}_4 + \left(\frac{28}{3}c_1^2 - 3c_2\right)\widetilde{\mathcal{A}}_5 + \cdots \ , \tag{29}$$

$$\mathcal{A}_4 = \widetilde{\mathcal{A}}_4 - \frac{13}{3}c_1\widetilde{\mathcal{A}}_5 + \cdots \ , \qquad \mathcal{A}_5 = \widetilde{\mathcal{A}}_5 + \cdots \ , \qquad \text{etc.} \tag{30}$$

In Ref. [41] this analytization was extended to the case when $k \mapsto \nu$ is noninteger[13]

$$\widetilde{\mathcal{A}}_{\nu+1}(Q^2) = \frac{1}{\pi}\frac{(-1)}{\beta_0^\nu \Gamma(\nu+1)}\int_0^\infty \frac{d\sigma}{\sigma}\rho_1(\sigma)\mathrm{Li}_{-\nu}\left(-\frac{\sigma}{Q^2}\right) \quad (-1 < \nu) \ , \tag{31}$$

where, as always, $\rho_1(\sigma) \equiv \mathrm{Im}\ \mathcal{A}_1(Q^2 = -\sigma - i\epsilon)$, and $\mathrm{Li}_{-\nu}(z)$ is the polylogarithm function.[14] The corresponding analogs of powers $a_{\rm pt}^\nu$ are then obtained by using the general relations

$$a_{\rm pt}^\nu = \widetilde{a}_{{\rm pt},\nu} + \sum_{m=1}^{\infty} \widetilde{k}_m(\nu)\, \widetilde{a}_{{\rm pt},\nu+m} \ . \tag{32}$$

and the linearity of analytization, i.e.

$$\mathcal{A}_\nu(Q^2) \equiv \left(a_{\rm pt}^\nu(Q^2)\right)_{\rm an} = \widetilde{\mathcal{A}}_\nu(Q^2) + \sum_{m\geq 1}\widetilde{k}_m(\nu)\,\widetilde{\mathcal{A}}_{\nu+m}(Q^2) \quad (\nu > -1) \ . \tag{33}$$

The coefficients $\widetilde{k}_m(\nu)$, for general real ν and positive integer m, were calculated in Ref. [41], and are combinations of Gamma functions and its derivatives, with arguments involving $\nu + \ell$ (ℓ being various integers).

Furthermore, since $a^\nu \ln^k a = \partial^k a^\nu / \partial \nu^k$, the linearity of analytization then implies

$$\begin{aligned}\mathcal{A}_{\nu,k}(Q^2) \equiv \left(a_{\rm pt}^\nu(Q^2)\ln^k a_{\rm pt}(Q^2)\right)_{\rm an} &= \frac{\partial^k \mathcal{A}_\nu(Q^2)}{\partial \nu^k} \\ &= \frac{\partial^k \widetilde{\mathcal{A}}_\nu(Q^2)}{\partial \nu^k} + \sum_{m\geq 1}\sum_{s=0}^{k}\binom{k}{s}\frac{\partial^s \widetilde{k}_m(\nu)}{\partial \nu^s}\,\frac{\partial^{k-s}\widetilde{\mathcal{A}}_{\nu+m}(Q^2)}{\partial \nu^{k-s}} \quad (\nu > -1) \ ,\end{aligned} \tag{34}$$

where in the terms on the right-hand side we use expressions for $\widetilde{\mathcal{A}}_{\nu+m}(Q^2)$ obtained by Eq. (31). Comparing Eqs. (34) and (31), we see that the terms in the above sum represent integrals over the scale σ involving the basic discontinuity function of the model (ρ_1) and derivatives of the polylogarithm function $\mathrm{Li}_{-\nu-m+1}$ with respect to its index ν. In the evaluation of the binding energy $E_{q\bar{q}}$ we will encounter the logarithmic terms of the type (34) with ν integer ($\nu = n$); however, the derivatives with respect to index ν in Eq. (34) imply that we must know the behavior of $\widetilde{\mathcal{A}}_\nu$ around the integer value $\nu = n$, i.e., we need to use here the expressions (31) for noninteger ν, or a version of it with improved integration convergence, Eq. (22) of Ref. [41] (cf. also the earlier footnote 13).

The analytization of expansions involving powers $a_{\rm pt}^\nu(\mu^2)$ and $a_{\rm pt}^\nu(\mu^2)\ln^k a_{\rm pt}(\mu^2)$ now consists simply in the replacements

$$\begin{aligned} a_{\rm pt}^\nu(\mu^2) &\mapsto \left(a_{\rm pt}^\nu(\mu^2)\right)_{\rm an} = \mathcal{A}_\nu(\mu^2) \ , \\ a_{\rm pt}^\nu(\mu^2)\ln^k a_{\rm pt}(\mu^2) &\mapsto \left(a_{\rm pt}^\nu(\mu^2)\ln^k a_{\rm pt}(\mu^2)\right)_{\rm an} = \mathcal{A}_{\nu,k}(\mu^2) \ .\end{aligned} \tag{35}$$

In the special case of MA/APT, this analytization is, in principle, equivalent to that of Eqs. (2)-(3), as argued in Refs. [21, 41]. The (small) differences between the two analytizations in MA/APT arise due to somewhat different truncations applied in the two approaches. One is the truncation in the loop-expansion ($\leq a_{\rm pt}^{\ell+1}$) usually applied to the running coupling $a_{\rm pt}(\mu^2)$, which is then reflected in the spectral functions $\mathrm{Im}a_{\rm pt}^\nu(-\sigma - i\epsilon)$ and $\mathrm{Im}[a_{\rm pt}^\nu(-\sigma -$

---

[13] This relation was also reformulated so as to be applicable in a larger ν interval: $-2 < \nu$, cf.Eq. (22) of Ref. [41].

[14] In Mathematica [48] it is implemented under the name PolyLog$[-\nu, z]$.



FIG. 1: The analytic couplings $\mathcal{A}_1(\mu^2)$ at low positive μ^2 ($\equiv Q^2$), for MA/APT and 2danQCD. Included are also pQCD couplings $a_{\rm pt}(\mu^2)$ in $\overline{\rm MS}$ and Lambert scheme. See the text for details.



FIG. 2: (a) The squares of the pQCD couplings, $a_{\rm pt}^2$, and their analytic analogs $\mathcal{A}_2$, at low positive μ^2; (b) the logarithmic derivatives $\widetilde{a}_{{\rm pt},2}$ and $\widetilde{\mathcal{A}}_2$ at low positive μ^2. See the text for details.

$i\epsilon)\ln^k a_{\rm pt}(-\sigma-i\epsilon)]$ in Eqs. (2)-(3). In the general approach, Eqs. (31)-(34), which can be applied also to the MA/APT analytic model, the same kind of truncation for $a_{\rm pt}(\mu^2)$ can be applied in $\rho_1^{(\rm pt)}(\sigma) = {\rm Im}\ a_{\rm pt}(Q^2=-\sigma-i\epsilon)$ in the integral (31), and the corresponding loop-truncation $m \leq m_{\rm max} = \ell - \nu + 1$ in the summations (33) and (34).

In Fig. 1 we present the analytic running couplings $\mathcal{A}_1(\mu^2)$ of 2danQCD and MA/APT, at positive low values of μ^2 ($\equiv Q^2$), for the central input values. In MA/APT this means: $\Lambda_{n_f=3} = 0.948$ GeV, $c_2 = c_2(\overline{\rm MS}, N_f=3) \approx +4.471$, $c_j = c_2^{j-1}/c_1^{j-2}$ for $j \geq 3$; in 2danQCD: see the second line of Table I, $c_2 = -4.76$, $c_j = c_2^{j-1}/c_1^{j-2}$ for $j \geq 3$. For comparison, we also include the pQCD couplings $a_{\rm pt}(\mu^2)$, in the Lambert renormalization scheme (i.e., the scheme of 2danQCD), and in the 4-loop $\overline{\rm MS}$ scheme: $c_2 = c_2(\overline{\rm MS}, N_f=3) \approx +4.471$; $c_3 = c_3(\overline{\rm MS}, N_f=3) \approx +20.99$; $c_j = 0$ for $j \geq 4$. In Fig. 2(a) we present the higher couplings $\mathcal{A}_2$ and $a_{\rm pt}^2$, where in MA/APT the coupling $\mathcal{A}_2$ was constructed according to Eq. (2), and in 2danQCD according to Eq. (28) truncated at $\widetilde{\mathcal{A}}_5$ term [cf. also Eq. (27)]. While we can see in Fig. 1 that the Lambert $a_{\rm pt}$ and 2danQCD $\mathcal{A}_1$ practically agree with each other down to scales $\mu^2 \approx 1\ {\rm GeV}^2$, we can also see in Fig. 2(a) that Lambert $a_{\rm pt}^2$ differs more significantly from the constructed 2danQCD analog $\mathcal{A}_2$ even at somewhat higher scales, this difference being primarily the consequence of the aforementioned truncation at $\widetilde{\mathcal{A}}_5$ in the construction of $\mathcal{A}_2$. In Fig. 2(b) we present, for comparison, the logarithmic derivatives $\widetilde{a}_{{\rm pt},2}(\mu^2)$ and $\widetilde{\mathcal{A}}_2(\mu^2)$ [cf. Eqs. 20) and (27)], where we can see that the difference between these couplings in 2danQCD and the corresponding (Lambert) pQCD is again negligible for $\mu^2 > 1\ {\rm GeV}^2$.

The 2danQCD model has been successfully applied [33] in the analysis of $(V+A)$ τ-decay data via Borel sum rules with OPE.

III. PERTURBATION EXPANSION FOR HEAVY $q\bar{q}$ GROUND STATE ENERGY

The analysis of nonrelativistic potential is the starting point for the determination of the ground state energy of $\bar{q}q$ and thus of the mass of such systems. The main input in these calculations is the mass $\overline{m}_q(\overline{m}_q^2)$ (here denoted simply as $\overline{m}_q$), also called $\overline{\rm MS}$ quark mass. The masses of the heavy quarkonia $\bar{q}q$ ($\Upsilon(1S)$ when $q=b$, $J/\psi(1S)$ when $q=c$) are well measured, and this allows us to extract the corresponding mass $\overline{m}_q$. By evaluating an observable, such as the quark-antiquark binding energy here, within anQCD models, at least part of the (chirality-conserving) nonperturbative effects gets included in the leading-twist term via the analytization, such as Eqs. (2)-(3) in MA/APT, and (35) in general analytic QCD models.

The coefficients in the (leading-twist) perturbation expansion of the ground state binding energy $E_{q\bar{q}}$ (i.e., with: $n=1$ and $\ell=0$) of heavy quarkonium $q\bar{q}$ in powers of $a_{\rm pt}$ were obtained up to all terms $\mathcal{O}(a_{\rm pt}^4)$ in Ref. [58], the terms $\mathcal{O}(a_{\rm pt}^5 \ln a_{\rm pt})$ in Ref. [59], and all terms up to $\mathcal{O}(a_{\rm pt}^5)$ (including logarithmic) are given in Ref. [7]. The last term ($\sim a_{\rm pt}^5$) is now completely known since the parameter a_3 from the static potential is now known, Refs. [5, 6]. The general structure of the (leading-twist term of the) ground state binding energy in pQCD is

$$\begin{aligned} E_{q\bar{q}}^{(\rm pt)} &= -\frac{4}{9} m_q \pi^2 a_{\rm pt}^2(\mu^2) \Big\{ 1 + a_{\rm pt}(\mu^2) \left[K_{1,0} + K_{1,1} L_{\rm pt}(\mu^2) \right] + a_{\rm pt}^2(\mu^2) \left[K_{2,0} + K_{2,1} L_{\rm pt}(\mu^2) + K_{2,2} L_{\rm pt}^2(\mu^2) \right] \\ & \quad + a_{\rm pt}^3(\mu^2) \left[K_{3,0,0} + K_{3,0,1} \ln a_{\rm pt}(\mu^2) + K_{3,1} L_{\rm pt}(\mu^2) + K_{3,2} L_{\rm pt}^2(\mu^2) + K_{3,3} L_{\rm pt}^3(\mu^2) \right] + \mathcal{O}(a_{\rm pt}^4) \Big\} \ , \end{aligned} \tag{36}$$

where

$$L_{\rm pt}(\mu^2; m_q^2) = \frac{1}{2} \ln \left(\frac{\mu^2}{((4\pi/3) m_q)^2 a_{\rm pt}^2(\mu^2)} \right) \ , \tag{37}$$

μ^2 is the (square of the) renormalization scale, m_q is the pole mass of the quark, and the coefficients $K_{j,k}$ can be obtained by combining the results of the aforementioned literature, presented also in Appendix A. The typical scale of the process is a so called soft scale Q_s^2 ($\equiv -q^2$) which represents a typical quark-antiquark momentum transfer inside the quarkonium ($Q_s^2 \sim m_q^2 \alpha_s^2$). The "soft" renormalization scale $\mu^2 \equiv \mu_s^2$ is expected to be of the same order

$$\mu^2 \equiv \mu_s^2 = \kappa Q_s^2 \quad (\kappa \sim 1; Q_s^2 \sim m_q^2 \alpha_s^2) \ . \tag{38}$$

The quarkonium mass is then

$$M_{q\bar{q}} = 2 m_q + E_{q\bar{q}}(m_q) \ . \tag{39}$$

In principle, the input quantity here could be the quark pole mass m_q. However, this mass m_q, in contrast to the mass $\overline{m}_q$, suffers from the strong infrared renormalon ambiguity (at Borel parameter value $b=1/2$, $\Rightarrow \delta m_q \sim \Lambda_{\rm QCD}$). This ambiguity must cancel in the physical sum, Eq. (39).

Therefore, it is more convenient to use as input a renormalon ambiguity-free input mass, such as $\overline{m}_q$, and we will follow this approach.

In the case of the bottom quark, before we relate the pole mass m_b with the mass $\overline{m}_b$, at renormalization energies $\mu = \overline{m}_b$ where $N_f = 4$ (and where the charm quark mass is considered zero, i.e., decoupled), the effects δm_b of the nonzero massa $m_c \neq 0$ have to be subtracted, and they are [60]

$$(\delta m_b)_{m_c} = (\delta m_b)_{m_c}^{(1)} + (\delta m_b)_{m_c}^{(2)} \approx 0.025 \pm 0.005 \ {\rm GeV} \ . \tag{40}$$

These effects were calculated in Ref. [60] in pQCD, at the hard renormalization scale $\mu^2 = \overline{m}_b^2$. We checked that they do not get significantly modified in MA/APT and in 2danQCD.

The pole mass m_q and the mass $\overline{m}_q$ are then related via the relation

$$\frac{m_q - \delta m_q}{\overline{m}_q} = 1 + R_0 \left(a_{\rm pt}(\overline{m}_q^2) + r_1 a_{\rm pt}^2(\overline{m}_q^2) + r_2 a_{\rm pt}^3(\overline{m}_q^2) + r_3 a_{\rm pt}^4(\overline{m}_q^2) \right) + \mathcal{O}(a_{\rm pt}^5) \ , \tag{41}$$

where δm_q is zero when $q=c$, and is given by Eq. (40) when $q=b$

$$\delta m_q \equiv \left\{ \begin{array}{ll} (\delta m_b)_{m_c} & (q=b) \\ 0 & (q=c) \end{array} \right\} \ . \tag{42}$$

The coefficients R_0, r_1, and r_2 are known (Refs. [61], [62], [63–65], respectively)

$$R_0 = \frac{4}{3} \ , \quad R_0 r_1 \ = \ 6.248\beta_0 - 3.739 \ , \tag{43}$$

$$R_0 r_2 \ = \ 23.497\beta_0^2 + 6.248\beta_1 + 1.019\beta_0 - 29.94 \ , \tag{44}$$

where $\beta_0 = (11 - 2N_f/3)/4$ and $\beta_1 = (102 - 38N_f/3)/16$ are the two universal aforementioned RGE coefficients, and N_f is the number of flavors of quarks lighter than q. Specifically, the values are: $r_1 = 7.739$ and $r_2 = 87.224$ for $N_f = 3$; and $r_1 = 6.958$ and $r_2 = 70.659$ for $N_f = 4$. On the other hand, in Appendix B we estimate the values of r_3 to a reasonably high level of precision (with less than 4 % uncertainty) by a method which uses the structure of the leading infrared renormalon (at $b = 1/2$) of the quantity $m_q/\overline{m}_q$

$$r_3(N_f = 3) = 1339.4 \ , \quad r_3(N_f = 4) = 987.3 \ . \tag{45}$$

While the relation (41) is written here at the "hard" renormalization scale $\mu^2 = \overline{m}_q^2$, it is straightforward to reexpress the sum on the right-hand side of Eq. (41) at a different, lower, scale μ^2. In Appendix B this reexpression is presented explicitly, under the assumption that during the lowering of the scale, $\overline{m}_q^2 \to \mu^2$, we do not cross quark threshold. The goal is to express in the perturbation expansion of the binding energy (36), where the renormalization scale is soft, the pole mass m_q via the mass $\overline{m}_q$, and for this we need the relation (41) at the soft renormalization scale. It turns out that for the $b\bar{b}$ system the hard scale is $\overline{m}_b \approx 4$ GeV, i.e., the scale where $N_f = 4$; and the soft scale, Eq. (38), is $\mu_s \approx 2$ GeV, i.e., the scale where it is more reasonable to expect $N_f = 3$.[15] In this case, we take into account also the (three-loop) quark threshold transition $N_f = 4 \mapsto 3$ at $\mu^2 = (2m_c)^2$, Ref. [66]. We thus obtain the relation (41) reexpressed at the soft scale μ_s^2 of Eq. (38)

$$m_q(\overline{m}_b^2;\mu_s^2) = \delta m_q + \overline{m}_q \left\{ 1 + \frac{4}{3} a_{\rm pt}(\mu_s^2) \left[1 + a_{\rm pt}(\mu_s^2) r_1(\mu_s^2) + a_{\rm pt}^2(\mu_s^2) r_2(\mu_s^2) + a_{\rm pt}^3(\mu_s^2) r_3(\mu_s^2) \right] + \mathcal{O}(a_{\rm pt}^5) \right\} \ . \tag{46}$$

Further, the renormalization scheme can also be varied in this relation, i.e., the changes of the scheme parameters $c_j = \beta_j/\beta_0$ $(j = 2,3)$ from the usual $\overline{\rm MS}$ scheme to other schemes affect correspondingly the values of the coupling $a_{\rm pt}(\mu_s^2)$ and of the coefficients $r_k(\mu_s^2)$. The relationship between $a_{\rm pt}$'s at two different scales and in two different renormalization schemes is summarized in Appendix C, where we also summarize the (3-loop) connection of $a_{\rm pt}$'s across the quark threshold.

After carefully performing all these transformations, we can rewrite the original expansion (36) for the binding energy in terms of the $\overline{m}_q$ quark mass, at any soft renormalization scale μ_s and in any chosen renormalization scheme $(c_2, c_3, \ldots)$

$$\begin{aligned} E_{q\bar{q}}^{(\rm pt)}(Q_s^2;\overline{m}_q^2;N_f=3) = & -\frac{4}{9}(\overline{m}_q + \delta m_q)\pi^2 a_{\rm pt}^2(\mu_s^2) \Big\{ 1 + a_{\rm pt}(\mu_s^2) \left[\mathcal{K}_{1,0} + \mathcal{K}_{1,1}\mathcal{L}_{\rm pt}(\mu_s^2) \right] \\ & + a_{\rm pt}^2(\mu_s^2) \left[\mathcal{K}_{2,0} + \mathcal{K}_{2,1}\mathcal{L}_{\rm pt}(\mu_s^2) + \mathcal{K}_{2,2}\mathcal{L}_{\rm pt}^2(\mu_s^2) \right] \\ & + a_{\rm pt}^3(\mu_s^2) \left[\mathcal{K}_{3,0,0} + \mathcal{K}_{3,0,1} \ln a_{\rm pt}(\mu_s^2) + \mathcal{K}_{3,1}\mathcal{L}_{\rm pt}(\mu_s^2) + \mathcal{K}_{3,2}\mathcal{L}_{\rm pt}^2(\mu_s^2) + \mathcal{K}_{3,3}\mathcal{L}_{\rm pt}^3(\mu_s^2) \right] + \mathcal{O}(a_{\rm pt}^4) \Big\} \ , \end{aligned} \tag{47}$$

where the logarithm contains now $\overline{m}_q$ quark mass

$$\mathcal{L}_{\rm pt}(\mu_s^2;\overline{m}_q^2) \ = \ \frac{1}{2} \ln \left(\frac{\mu_s^2}{\left[(4\pi/3)(\overline{m}_q + \delta m_q) \right]^2 a_{\rm pt}^2(\mu_s^2)} \right) \ . \tag{48}$$

We note that the new (renormalon-ambiguity-free) mass which appears naturally in this expansion is not exactly the mass $\overline{m}_q$, but rather

$$\widetilde{m}_q \equiv \overline{m}_q + \delta m_q = \left\{ \begin{array}{ll} \overline{m}_b + (\delta m_b)_{m_c} & (q = b) \\ \overline{m}_c & (q = c) \end{array} \right\} \ , \tag{49}$$

where $(\delta m_b)_{m_c}$ is given by Eq. (40) when $q = b$.

---

[15] Usually, the quark thresholds are taken at $2m_q$. In the case of $N_f = 4 \mapsto 3$ transition, this is about 2.5 GeV, above the soft scale of the $b\bar{b}$ system.

The aforementioned soft "process scale" Q_s^2 ($\sim \overline{m}_q^2\alpha_s^2$) can be regarded, at least formally, to be a variable complex scale. Therefore, the binding energy $E_{q\bar{q}}(Q_s^2;\overline{m}_q^2)$ is, formally, a spacelike observable analytic in Q_s^2. The analytization of $E_{q\bar{q}}^{(\mathrm{pt})}(Q_s^2;\overline{m}_q^2)$ of Eq. (47), according to Eqs. (35) [or, in MA/APT: Eqs. (2)-(3)], then leads to

$$
\begin{aligned}
E_{q\bar{q}}^{(\mathrm{an})}(Q_s^2;\overline{m}_q^2) = -\frac{4}{9}\widetilde{m}_q\pi^2\Big\{ & \mathcal{A}_2(\kappa Q_s^2) + \left[\mathcal{K}_{1,0}\mathcal{A}_3(\kappa Q_s^2) + \mathcal{K}_{1,1}\mathcal{B}_{3,1}(\kappa Q_s^2)\right] \\
& + \left[\mathcal{K}_{2,0}\mathcal{A}_4(\kappa Q_s^2) + \mathcal{K}_{2,1}\mathcal{B}_{4,1}(\kappa Q_s^2) + \mathcal{K}_{2,2}\mathcal{B}_{4,2}(\kappa Q_s^2)\right] \\
& + \left[\mathcal{K}_{3,0,0}\mathcal{A}_5(\kappa Q_s^2) + \mathcal{K}_{3,0,1}\mathcal{A}_{5,1}(\kappa Q_s^2) + \mathcal{K}_{3,1}\mathcal{B}_{5,1}(\kappa Q_s^2) + \mathcal{K}_{3,2}\mathcal{B}_{5,2}(\kappa Q_s^2) + \mathcal{K}_{3,3}\mathcal{B}_{5,3}(\kappa Q_s^2)\right] + \mathcal{O}(\mathcal{A}_{6,4})\Big\} \ , \qquad (50)
\end{aligned}
$$

where we use, for simplicity, the notation of Eq. (33) with Eq. (27) for $\mathcal{A}_\nu$'s ($\nu = k$ integer now) [in MA/APT: Eq. (2)], and denote by $\mathcal{B}_{n+2,j}$ the following:

$$
\begin{aligned}
\mathcal{B}_{n+2,j}(\kappa Q_s^2) &= \left(a_{\mathrm{pt}}^{n+2}(\kappa Q_s^2)\frac{1}{2^j}\ln^j\left(\frac{\kappa Q_s^2}{((4\pi/3)\overline{m}_q)^2 a_{\mathrm{pt}}^2(\kappa Q_s^2)}\right)\right)_{\mathrm{an}} \\
&= \frac{1}{2^j}\sum_{s=0}^{j}\binom{j}{s}(-2)^s\ \bar{f}^{j-s}(\kappa Q_s^2)\mathcal{A}_{n+2,s}(\kappa Q_s^2) \ , \qquad (51)
\end{aligned}
$$

where

$$
\bar{f} \equiv \ln\left(\frac{\kappa Q_s^2}{(4\pi/3)^2\overline{m}_q^2}\right) \ , \qquad (52)
$$

and $\mathcal{A}_{n+2,s}$ were defined in Eq. (34) [in MA/APT: Eq. (3)].

Here, we have $j = 1, \ldots, n$, implying that $\mathcal{B}_{n+2,j}(\kappa Q_s^2) \to 0$ faster than $a_{\mathrm{pt}}^2(\kappa Q_s^2)$ when $|Q_s^2| \to \infty$, by asymptotic freedom.

The relation (46) in its analytized form is

$$
m_q(\overline{m}_b^2;\mu_s^2) = \delta m_q + \overline{m}_q\left\{1 + \frac{4}{3}\left[\mathcal{A}_1(\mu_s^2) + r_1(\mu_s^2)\mathcal{A}_2(\mu_s^2) + r_2(\mu_s^2)\mathcal{A}_3(\mu_s^2) + r_3(\mu_s^2)\mathcal{A}_4(\mu_s^2)\right] + \mathcal{O}(\mathcal{A}_5)\right\} \ . \qquad (53)
$$

Since this relation includes at its highest order the term $\sim \mathcal{A}_4$ ($\sim a_{\mathrm{pt}}^4$), in general analytic QCD it is consistent to use in Eq. (53) for the expressions $\mathcal{A}_n$ ($n = 1,2,3,4$) those given in Eq. (33) [or: Eqs. (28)-(30)] with the sum there truncated at (and including) $\widetilde{\mathcal{A}}_4$ ($\sim \mathcal{A}_4 \sim a_{\mathrm{pt}}^4$). On the other hand, the expression (50) for the binding energy includes at its highest order the terms $\sim \mathcal{A}_5$ (more precisely: $\sim \mathcal{A}_{5,3}$), therefore it is consistent to use there for the expressions $\mathcal{A}_n$ ($2 \le n \le 5$) those of Eq. (33) [or: Eqs. (28)-(30)] with the sum truncated at (and including) $\widetilde{\mathcal{A}}_5$, and expressions $\widetilde{\mathcal{A}}_{n,k}$ of Eq. (34) truncated at (and including) $\partial^k\widetilde{\mathcal{A}}_{\nu+5}/\partial\nu^k|_{\nu=0}$ ($\sim a_{\mathrm{pt}}^n \ln^k a_{\mathrm{pt}}$), where $k = 1,2,3$.

The aforementioned soft scale Q_s^2 ($\sim \overline{m}_q^2\alpha_s^2$) can be fixed in pQCD, by convention, by the condition that all the logarithmic terms in the expansion (47) disappear

$$
Q_{s,\mathrm{pt}}^2 = (4\pi/3)^2\widetilde{m}_q^2 a_{\mathrm{pt}}^2(Q_{s,\mathrm{pt}}^2) \ , \qquad (54)
$$

where $\widetilde{m}_q$ is defined in Eq. (49). This type of condition, in analytic QCD, would correspond to fixing a referential soft scale Q_s by requiring $\mathcal{B}_{n+1,j}(Q_s^2) = 0$, for various n's and $j = 1,\ldots,n$, meaning that the fixing is not unique since it depends on n and j. Our convention will be to require that the leading logarithmic term $\mathcal{B}_{n+1,j}$ in Eq. (50), i.e., $\mathcal{B}_{3,1}$, be zero at such scale

$$
\mathcal{B}_{3,1}(Q_s^2) = 0 \ . \qquad (55)
$$

It will turn out that this condition has a solution in the case of $b\bar{b}$ ($\Upsilon(1S)$) in 2danQCD, but not in $J/\psi(1S)$ in that model, and not in any case of the MA/APT model. In such respective cases, we will use simply the following, more naive, analogs of the pQCD condition (54):

$$
\begin{aligned}
Q_s^2 &= (4\pi/3)^2\widetilde{m}_q^2\widetilde{\mathcal{A}}_2(Q_s^2) \quad (J/\psi(1S) \text{ in 2danQCD}) \ , \qquad (56) \\
Q_{s,(\mathrm{MA})}^2 &= (4\pi/3)^2\widetilde{m}_q^2\mathcal{A}_2^{(\mathrm{MA})}(Q_{s,(\mathrm{MA})}^2) \quad (\text{MA/APT}) \ . \qquad (57)
\end{aligned}
$$

These measures of the typical momentum scale of the (nonrelativistic) quark inside the quarkonium are rather low, ≈ 2 GeV in $\Upsilon(1S)$, and $\lesssim 1$ GeV in $J/\psi(1S)$. In pQCD such scales are problematic, because they are not far away from the unphysical (Landau) singularities of $a_{\rm pt}(Q^2)$; in ananlytic QCD models, no such problems appear in principle.

We recall that the pole mass m_q and the static potential $V(r)$ both contain the leading infrared renormalon ($b=1/2$) singularities, and that cancellation of these singularities takes place in the sum $2m_q+V(r)$ [67–69]. As a consequence, this cancellation must take place also in the quarkonium mass $2m_q+E_{q\bar q}$ [60, 70–74], more specifically, in the sum $2m_q+E_{q\bar q}(s)$ where $E_{q\bar q}(s)=\langle 1|V(r)|1\rangle$ ($\sim V(r_s)$) is the soft part of the binding energy $E_{q\bar q}$, and $|1\rangle$ denotes the (ground) state of the quarkonium. The typical soft distance r_s in the quarkonium is $\sim 1/\sqrt{Q_s^2}$ where Q_s^2 is the soft scale of Eqs. (55)-(57), i.e. $1/r_s \sim (m_q \pi a)$ (where $a=a_{\rm pt}$ or $\mathcal{A}_1$). However, since $V(r_s) \propto 1/r_s$, we have $E_{q\bar q}(s)/m_q \sim V(r_s)/m_q \propto 1/(r_s m_q) \sim (\pi a)$, this thus leading to the so called "power mismatch" in the renormalon cancellation in the pQCD expansions $2m_q+E_{q\bar q}(s)$ [75] (see also: [70]): the terms $\sim a^n$ in $2m_q/\overline{m}_q$ tend to cancel numerically the terms $\sim a^{n+1}$ in $E_{q\bar q}/\overline{m}_q$. Therefore, since the binding energy $E_{q\bar q}$ is now known up to $\sim a_{\rm pt}^5$, it is very convenient to have the relation $m_q/\overline{m}_q$ up to $\sim a^4$, i.e., to have a good estimate for the coefficient r_3 and to use it, so that the effects of renormalon cancellation in $2m_q+E_{q\bar q}(s)$ can be seen numerically more clearly. Of course, this cancellation, term-by-term, should be numerically more precise (at sufficiently high orders) if the renormalization scales μ_s used in $m_q/\overline{m}_q$ and in $E_{q\bar q}(s)/\overline{m}_q$, Eqs. (46) and (47), are taken to be equal. This renormalon cancellation will be our guiding principle for the separation of the ultrasoft (us) part from the binding energy

$$E_{q\bar q} = E_{q\bar q}(s) + E_{q\bar q}(us) \ . \tag{58}$$

Typical us scales are $\mu_{us} \sim \overline{m}_q \alpha_s^2$, and the us part of the binding energy is $\sim \overline{m}_q a^5 \ln a$. We can parametrize the s-us separation by a dimensionless parameter $k_{s/us}$ such that the s-us factorization scale μ_f is written as

$$\mu_f = k_{s/us} \overline{m}_q \alpha_s(Q_s)^{3/2} \left[\approx k_{s/us} (Q_s Q_{us})^{1/2} \right] \ , \tag{59}$$

It is expected that usually $k_{s/us} \sim 1$, but it does not have to be so always. The us part can be rewritten, in terms of $k_{s/us}$ as (cf. Ref. [74])

$$E_{q\bar q}^{\rm (pt)}(us) \ = \ -\frac{4}{9} \widetilde{m}_q \pi^2 \left[\mathcal{K}_{3,0,0}(us) a_{\rm pt}^5(\mu_{us}^2) + \mathcal{K}_{3,0,1}(us) a_{\rm pt}^5(\mu_{us}^2) \ln a_{\rm pt}(\mu_{us}^2) + \mathcal{O}(a_{\rm pt}^6) \right] \ , \tag{60}$$

and in analytic QCD correspondingly

$$E_{q\bar q}^{\rm (an)}(us) \ = \ -\frac{4}{9} \widetilde{m}_q \pi^2 \left[\mathcal{K}_{3,0,0}(us) \mathcal{A}_5(\mu_{us}^2) + \mathcal{K}_{3,0,1}(us) \mathcal{A}_{5,1}(\mu_{us}^2) + \mathcal{O}(\mathcal{A}_6) \right] \ , \tag{61}$$

where μ_{us} is a us renormalization scale ($\mu_{us} \sim \overline{m}_q \alpha_s^2$), and the two us coefficients are

$$\mathcal{K}_{3,0,0}(us) \ = \ \left[27.512 + 7.098 \ln \pi - 14.196 \ln(k_{s/us}) \right] \pi^3 \ , \tag{62}$$

$$\mathcal{K}_{3,0,1}(us) \ = \ 7.098 \pi^3 \ . \tag{63}$$

The expansion of the soft (s) part $E_{q\bar q}(s)$ of the binding energy is then, according to Eq. (58), the same as the expansions (47) and (50), with the exception of the replacements[16] of two coefficients $\mathcal{K}_{3,0,0}$ and $\mathcal{K}_{3,0,1}$

$$E_{q\bar q}^{\rm (pt,an)}(s) \ = \ E_{q\bar q}^{\rm (pt,an)} \left(\mathcal{K}_{3,0,0} \mapsto \mathcal{K}_{3,0,0} - \mathcal{K}_{3,0,0}(us); \mathcal{K}_{3,0,1} \mapsto \mathcal{K}_{3,0,1} - \mathcal{K}_{3,0,1}(us) \right) \ . \tag{64}$$

The s-us factorization, i.e., the parameter $k_{s/us}$, will then be determined, in each model, by requiring that the leading infrared renormalon cancellation in $2m_q+E_{q\bar q}(s)$ be exact at the last available order, i.e., that the $\mathcal{O}(a^4)$ term in $2m_q(Q_s^2, \overline{m}_q^2)$ and the term $\mathcal{O}(a^5)$[17] in $E_{q\bar q}(s; Q_s^2; \overline{m}_q^2)$ cancel exactly.

The us part of the quarkonium mass, $E_{q\bar q}(us; Q_{us}^2; \overline{m}_q^2)$, will be evaluated in each case according to a procedure which takes into account those problems of low-scale evaluations which appear in the considered model (pQCD, MA/APT, 2danQCD).

---

[16] The coefficients $\mathcal{K}_{3,0,0}(us)$ and $\mathcal{K}_{3,0,1}(us)$, representing the leading part of the (quasi)observable $E_{q\bar q}(us)$, are renormalization scale (μ_{us}) and scheme independent.

[17] The latter term includes all $\mathcal{O}(a^5 \ln^k a)$ terms ($k=0,1,2,3$).

IV. NUMERICAL RESULTS

A. Bottom mass extraction

In this Section, we extract from the mass of $b\bar{b}$ quarkonium the mass $\overline{m}_b \equiv \overline{m}_b(\overline{m}_b^2)$ in MA/APT, 2danQCD model, and in pQCD in two renormalization schemes ($\overline{\rm MS}$ and in the Lambert scheme of 2danQCD). For this, we use the relation between $\overline{m}_b$ and the well-measured mass of the $b\bar{b}$ quarkonium $\Upsilon(1S)$ [44]

$$2m_b(\overline{m}_b^2;\mu_s^2) + E_{b\bar{b}}(s;Q_s^2;\overline{m}_b^2;\mu_s^2,N_f=3) + E_{b\bar{b}}(us;Q_{us}^2;\overline{m}_b^2;\mu_{us}^2) = M_{\Upsilon(1S)}^{(\rm exp)}(=9.460 \ {\rm GeV}) \ . \tag{65}$$

The dependence on the (soft) renormalization scale μ_s^2 in the pole mass m_b and in the soft binding energy $E_{b\bar{b}}(s)$ occurs due to the truncation of the series for these two quantities. For the same reason, the ultrasoft binding energy $E_{b\bar{b}}(us)$ has strong dependence on the ultrasoft renormalization scale μ_{us}^2 due to the drastic truncation of this quantity at its leading order ($\sim a^5 \ln a$). As mentioned in the previous Section, the separation of the s and us parts of the binding energy will be performed here by determining the s-us separation parameter $k_{s/us}$, Eqs. (58)-(64), by the requirement of cancellation of the leading renormalon in $2m_b + E_{b\bar{b}}(s)$.

In contrast to the other three models, MA/APT gives a very small central value for the s-us separation parameter $k_{s/us} \approx 10.^{-12}$. This reflects the difficulty in the MA/APT scenario to exactly enforce the leading renormalon cancellation of the $\sim \mathcal{A}_4$ term of $2m_b$ with the corresponding $\sim \mathcal{A}_5$ term of $E_{b\bar{b}}(s)$. If, on the other hand, we impose in MA/APT the condition $k_{s/us} \sim 1$, more specifically the central value $k_{s/us} = 1$ and variation of that parameter in the interval $(0.1, 10.)$, the results change somewhat, the central extracted value of $\overline{m}_b$ increases by about 0.050 GeV, and the absolute values of the us part of the binding energy and of various other uncertainties of $\overline{m}_b$ get significantly reduced. We will consider in MA/APT only the natural range ~ 1 of the s-us separation parameter, $k_{s/us} = 1.0^{+9.0}_{-0.9}$, rather than the exceedingly small values of $k_{s/us}$ required by the exact renormalon cancellation. In the other three models (2danQCD, and in Lambert and $\overline{\rm MS}$ pQCD), the renormalon cancellation is imposed without any problems, resulting in the values of $k_{s/us}$ within the interval between 10^{-1} and 10^{1}.

In the previous Section we mentioned that we take $N_f = 3$ for the number of active flavors in the binding energy, i.e., in this case the m_c mass is considered to be infinite (decoupled). It turns out that, while the effects of the finiteness of m_c cannot be neglected in the relation between m_b and $\overline{m}_b$, Eqs. (40)-(41), these effects can be safely neglected in the binding energy $E_{b\bar{b}}(N_f=3)$, cf. Ref. [60] based on Refs. [62, 76, 77].

Application of the formalism described in Subsec. II A and Subsec. II B for the calculation of the couplings of the analytic QCD models MA/APT and 2danQCD, and in Sec. III for the calculation of $2m_b$, $E_{b\bar{b}}(s)$ and $E_{\bar{b}b}(us)$ in terms of these couplings, then gives us the following results:

$$\begin{aligned}
\overline{m}_b({\rm MA/APT}) &= \Bigg\{ 4.165 \pm 0.002_{(us)} + \begin{pmatrix} +0.005 \\ -0.004 \end{pmatrix}_{(s/us)} + \begin{pmatrix} -0.019 \\ +0.020 \end{pmatrix}_{(\Lambda^2)} \\
&\quad + \begin{pmatrix} -0.005 \\ +0.003 \end{pmatrix}_{(\mu_s^2)} \mp 0.005_{(m_c)} \Bigg\} \ {\rm GeV} && (66) \\
&= 4.165 \pm 0.022 \ {\rm GeV} \ , \quad {\rm with:} \ Q_{s,({\rm MA})} = 1.60 \ {\rm GeV} \ , \ k_{s/us} = 1.0 \ . && (67)
\end{aligned}$$

$$\begin{aligned}
\overline{m}_b({\rm 2danQCD}) &= \Bigg\{ 4.353 + \begin{pmatrix} -0.068 \\ +0.071 \end{pmatrix}_{(us)} + \begin{pmatrix} +0.015 \\ -0.016 \end{pmatrix}_{(s/us)} \mp 0.005_{(\alpha_s)} \\
&\quad + \begin{pmatrix} -0.023 \\ +0.034 \end{pmatrix}_{(c_2)} + \begin{pmatrix} +0.017 \\ -0.025 \end{pmatrix}_{(\mu_s^2)} \mp 0.005_{(m_c)} \Bigg\} \ {\rm GeV} && (68) \\
&= 4.353 \pm 0.084 \ {\rm GeV} \ , \quad {\rm with:} \ Q_s = 2.08 \ {\rm GeV} \ , \ k_{s/us} = 0.241 \ . && (69)
\end{aligned}$$

We will comment on the above uncertainties below. For completeness, we give here also the results of the same kind of analysis in pQCD, first in the Lambert renormalization scheme (i.e., the scheme as used in 2danQCD: $c_2 = -4.76$, $c_j = c_2^{j-1}/c_1^{j-2}$ for $j \geq 3$); and in (4-loop) $\overline{\rm MS}$ scheme:

$$\begin{aligned}
\overline{m}_b({\rm pQCDLamb.}) &= \Bigg\{ 4.382 + \begin{pmatrix} -0.091 \\ +0.097 \end{pmatrix}_{(us)} + \begin{pmatrix} -0.013 \\ +0.017 \end{pmatrix}_{(s/us)} \pm 0.010_{(\alpha_s)} \\
&\quad + \begin{pmatrix} +0.027 \\ -0.008 \end{pmatrix}_{(c_2)} + \begin{pmatrix} -0.002 \\ -0.041 \end{pmatrix}_{(\mu_s^2)} \mp 0.005_{(m_c)} \Bigg\} \ {\rm GeV} && (70) \\
&= 4.382 \pm 0.111 \ {\rm GeV} \ , \quad {\rm with:} \ Q_{s,{\rm pt}} = 1.73 \ {\rm GeV} \ , \ k_{s/us} = 0.306 \ . && (71)
\end{aligned}$$

$$\begin{aligned}
\overline{m}_b(\text{pQCD}\overline{\text{MS}}) &= \Big\{ 4.505 + \begin{pmatrix} -0.177 \\ +0.200 \end{pmatrix}_{(us)} + \begin{pmatrix} -0.082 \\ +0.084 \end{pmatrix}_{(s/us)} + \begin{pmatrix} +0.031 \\ -0.027 \end{pmatrix}_{(\alpha_s)} \\
&\quad + \begin{pmatrix} -0.004 \\ -0.075 \end{pmatrix}_{(\mu_s^2)} \mp 0.005_{(m_c)} \Big\} \ \text{GeV} \qquad (72) \\
&= 4.505 \pm 0.231 \ \text{GeV} \ , \quad \text{with} : Q_{s,\text{pt}} = 1.87 \ \text{GeV} \ , \ k_{s/us} = 0.248 \ . \qquad (73)
\end{aligned}$$

The value of the *s*-*us* separation parameter $k_{s/us}$ was determined in all cases by the aforementioned renormalon cancellation in the sum $M_{\Upsilon(1S)}(s) = 2m_b + E_{b\bar{b}}(s; k_{s/us})$, except in the MA/APT case, as discussed above. Below we present these resulting sums, for the central choices of the aforementioned four results, where we combine in each parenthesis the (positive) terms $\sim a^n$ of $2m_b$ and the corresponding (negative) terms $\sim a^{n+1}$ of $E_{b\bar{b}}(s)$ $(n = 0, 1, 2, 3, 4)$, in order to see more clearly the tendency of the renormalon cancellation; the *us* part is given separately

$$\begin{aligned}
M_{\Upsilon(1S)}(s; \text{MA/APT}) &= 8.380 + (1.151 - 0.152) + (0.281 - 0.173) + (0.137 - 0.152) + (0.075 - 0.092) \ \text{GeV} \\
&= 8.380 + 0.999 + 0.107 - 0.015 - 0.017 \ \text{GeV} \ (= 9.454 \ \text{GeV}) \ , \qquad (74) \\
E_{b\bar{b}}(us; \text{MA/APT}) &= 0.006 \mp 0.003 \ \text{GeV} \ , \qquad (\overline{m}_b = 4.165 \ \text{GeV}, \ k_{s/us} = 1.0) \ ; \qquad (75)
\end{aligned}$$

$$\begin{aligned}
M_{\Upsilon(1S)}(s; \text{2danQCD}) &= 8.756 + (0.999 - 0.131) + (0.373 - 0.222) + (0.233 - 0.193) + (0.568 - 0.568) \ \text{GeV} \\
&= 8.756 + 0.868 + 0.151 + 0.040 + 0.000 \ \text{GeV} \ (= 9.814 \ \text{GeV}) \ , \qquad (76) \\
E_{b\bar{b}}(us; \text{2danQCD}) &= -0.353 \pm 0.151 \ \text{GeV} \ , \qquad (\overline{m}_b = 4.353 \ \text{GeV}, \ k_{s/us} = 0.241) \ ; \qquad (77)
\end{aligned}$$

$$\begin{aligned}
M_{\Upsilon(1S)}(s; \text{pQCDLamb.}) &= 8.814 + (1.095 - 0.170) + (0.327 - 0.220) + (0.346 - 0.376) + (0.413 - 0.413) \ \text{GeV} \\
&= 8.814 + 0.925 + 0.107 - 0.031 + 0.000 \ \text{GeV} \ (= 9.816 \ \text{GeV}) \ , \qquad (78) \\
E_{b\bar{b}}(us; \text{pQCDLamb.}) &= -0.355 \pm 0.201 \ \text{GeV} \ , \qquad (\overline{m}_b = 4.382 \ \text{GeV}, \ k_{s/us} = 0.306) \ ; \qquad (79)
\end{aligned}$$

$$\begin{aligned}
M_{\Upsilon(1S)}(s; \text{pQCD}\overline{\text{MS}}) &= 9.060 + (1.183 - 0.193) + (0.399 - 0.263) + (0.330 - 0.439) + (0.475 - 0.475) \ \text{GeV} \\
&= 9.060 + 0.991 + 0.136 - 0.109 + 0.000 \ \text{GeV} \ (= 10.078 \ \text{GeV}) \ , \qquad (80) \\
E_{b\bar{b}}(us; \text{pQCD}\overline{\text{MS}}) &= -0.617 \pm 0.394 \ \text{GeV} \ , \qquad (\overline{m}_b = 4.505 \ \text{GeV}, \ k_{s/us} = 0.248) \ ; \qquad (81)
\end{aligned}$$

We can see from Eqs. (74)-(80) explicitly that for the chosen corresponding central values of the parameter $k_{s/us}$ the renormalon cancellation is exact in the last term (fifth term, named $t_5(s)$) in the sum for the soft mass $M_{\Upsilon(1S)}(s)$, except in the case of MA/APT where $k_{s/us} = 1.0$ was chosen and the cancellation in $t_5(s)$ is approximate.

In the extracted results for $\overline{m}_b$, Eqs. (66)-(73), the uncertainty from the ultrasoft (*us*) sector is rather strong, especially in the pQCD approaches. The *us* binding energy $E_{b\bar{b}}(us; \overline{m}_b^2; \mu_{us}^2)$, cf. Eqs. (59)-(63), depends strongly on the *s*-*us* separation parameter $k_{s/us}$, the central value of the latter being fixed by the renormalon cancellation of the soft mass $M_{\Upsilon(1S)}(s)$ in the last term (or, in MA/APT, $k_{s/us} = 1$). The renormalization scale μ_{us} was not allowed to go below $(\mu_{us})_{\min} = 1.1$ GeV, because at $\mu_{us} \approx 1.1$ GeV the coupling $\mathcal{A}_5(\mu_{us})$ $(= \widetilde{\mathcal{A}}_5(\mu_{us})$ in our approach$)$[18] in 2danQCD reaches a local maximum, indicating that the model may not necessarily give reliable values of $\mathcal{A}_5$ and $\mathcal{A}_{5,1}$ below such scales. Therefore, in general, we will not use scales lower than 1.1 GeV, in 2danQCD and in pQCD, i.e., in the *us* binding energy expressions (60)-(61). It turns out that the latter expression, Eq. (61), in 2danQCD reaches a local minimum at μ_{us} slightly above 1.1 GeV. Therefore, we estimate the *us* part of the binding energy in 2danQCD in the following way [we use the notation $E_{b\bar{b}}(us; \mu_{us}^2)$]:

$$E_{b\bar{b}}(us; \text{2danQCD}) = \frac{1}{2} \left[E_{b\bar{b}}(us; Q_s^2) + (E_{b\bar{b}}(us))_{\min} \right] \pm \frac{1}{2} \left[E_{b\bar{b}}(us; Q_s^2) - (E_{b\bar{b}}(us))_{\min} \right] \ , \qquad (82)$$

where the soft reference scale Q_s was determined by the condition (55), and gave for the central values of the input parameters (s_0 and c_2 of the second line of Table I; $\alpha_s(M_Z^2; \overline{\text{MS}}) = 0.1184$; $\overline{m}_b = 4.353$ GeV) the soft reference scale value $Q_s \approx 2.08$ GeV.

---

[18] We recall that in 2danQCD we calculate in $m_q(\overline{m}_q^2; \mu_s^2)$ the couplings $\mathcal{A}_n(\mu_s^2)$ as sums of $\widetilde{\mathcal{A}}_p(\mu_s^2)$ with $p = n, n+1, \ldots, 4$; and in $E_{q\bar{q}}(s; \overline{m}_b^2; \mu_s^2)$ and $E_{q\bar{q}}(us; \overline{m}_b^2; \mu_{us}^2)$ the couplings $\mathcal{A}_{n,k}(\mu_s^2)$ (and $\mathcal{A}_{n,k}(\mu_{us}^2)$) as (derivatives of the) sums of $\widetilde{\mathcal{A}}_p(\mu_s^2)$ (and $\widetilde{\mathcal{A}}_p(\mu_{us}^2)$) with $p = n, n+1, \ldots, 5$; cf. discussion in Subsec. II B and Sec. III.

In the two pQCD approaches, the us binding energy turns out to decrease monotonously when μ_{us} decreases below the soft reference scale $Q_{s,\mathrm{pt}}$ of Eq. (54). For pQCD in the Lambert scheme ($c_2 = -4.76$), with central values of the input parameters, the soft reference scale turns out to be $Q_{s,\mathrm{pt}} \approx 1.73$ GeV, and the us binding energy at $\mu_{us} = 1.1$ GeV reaches the value of -0.56 GeV; we use the following estimate for the us part of the binding energy:

$$E_{b\bar{b}}(us; \mathrm{Lamb.}) = \frac{1}{2}\left[E_{b\bar{b}}(us; Q^2_{s,\mathrm{pt}}) + E_{b\bar{b}}(us; 1.1^2 \mathrm{GeV}^2)\right] \pm \frac{1}{2}\left[E_{b\bar{b}}(us; Q^2_{s,\mathrm{pt}}) - E_{b\bar{b}}(us; 1.1^2 \mathrm{GeV}^2)\right] \ . \tag{83}$$

In pQCD in $\overline{\mathrm{MS}}$ scheme, however, $E_{b\bar{b}}(us; 1.1^2 \mathrm{GeV}^2) \approx -1.5$ GeV, which is exceedingly low and indicates failure of the method already at such scales, due to vicinity of Landau singularities of the running coupling.[19] Therefore, in $\overline{\mathrm{MS}}$ we take as the minimal acceptable scale $\mu_{us} = 1.2$ GeV, where $E_{b\bar{b}}(us; 1.2^2 \mathrm{GeV}^2) \approx -1.0$ GeV (and $Q_{s,\mathrm{pt}} \approx 1.87$ GeV) when the central values of the input parameters are used

$$E_{b\bar{b}}(us; \overline{\mathrm{MS}}) = \frac{1}{2}\left[E_{b\bar{b}}(us; Q^2_{s,\mathrm{pt}}) + E_{b\bar{b}}(us; 1.2^2 \mathrm{GeV}^2)\right] \pm \frac{1}{2}\left[E_{b\bar{b}}(us; Q^2_{s,\mathrm{pt}}) - E_{b\bar{b}}(us; 1.2^2 \mathrm{GeV}^2)\right] \ . \tag{84}$$

In the MA/APT case, on the other hand, Q_s is fixed according to Eq. (57) and gives, for the central input parameter values, the value $Q_{s,(\mathrm{MA})} \approx 1.60$ GeV. In MA/APT no practical problems appear at scales $\mu_{us} < 1.1$ GeV. It turns out that $E_{b\bar{b}}(us, \mathrm{MA/APT})$ reaches a moderate maximum value of 0.009 GeV at $\mu_{us} \approx 0.7$ GeV for the chosen central value $k_{s/us} = 1.0$. Therefore, we estimate the us part of the binding energy in MA/APT in the following way:

$$E_{b\bar{b}}(us; \mathrm{MA/APT}) = \frac{1}{2}\left[(E_{b\bar{b}}(us))_{\max} + E_{b\bar{b}}(us; Q^2_{s,(\mathrm{MA})})\right] \pm \frac{1}{2}\left[E_{b\bar{b}}(us; Q^2_{s,(\mathrm{MA})}) - (E_{b\bar{b}}(us))_{\max}\right] \ . \tag{85}$$

In Eqs. (66)-(73), the uncertainties in $\overline{m}_b$ originating from this determination of the us binding energy are denoted by the subscript (us).

The described behavior of the us binding energy $E_{b\bar{b}}(us; \mu^2_{us})$ in the three models (2danQCD, and pQCD in the two schemes), as a function of the us renormalization scales μ_{us} in the low-momentum regime, is presented in Fig. 3(a), and in the case of MA/APT in Fig. 3(b).



FIG. 3: (a) The ultrasoft binding energy $E_{b\bar{b}}(us)$ (in GeV) as a function of the us renormalization scale μ_{us}, for the central input values of 2danQCD model for the $\Upsilon(1S)$ system ($\overline{m}_b = 4.353$ GeV; $k_{s/us} = 0.241$; $c_2 = -4.76$; $s_0 = 23.06$): solid line for 2danQCD; dashed line for Lambert pQCD ($c_2 = -4.76$); dash-dotted line for the 4-loop $\overline{\mathrm{MS}}$ pQCD ($c_2 = 4.471$). For all three curves the same $\overline{m}_b$ and $k_{s/us}$ values are used. Included are also the properly rescaled 2danQCD couplings $\mathcal{A}_5$ and $\mathcal{A}_{5,1}$ as functions of the scale μ ($= \mu_{us}$). (b) The same as in Figure (a), but for the MA/APT model (with $c_2 = 4.471$), with the corresponding central values of that model ($\overline{m}_b = 4.165$ GeV and $k_{s/us} = 1$). In all the curves $N_f = 3$ was taken.

On the other hand, there are related uncertainties for the extracted $\overline{m}_b$, denoted by the subscript (s/us) in Eqs. (66)-(72), which originate from the variation of the s-us separation parameter $k_{s/us}$. The parameter $k_{s/us}$ was varied in such a way that the last (fifth, $t_5(s)$) term in the series for the soft mass $M_{\Upsilon(1S)}(s)$, cf. Eqs. (74)-(80), varies between the penultimate term $t_4(s)$ of these series, and its negative $-t_4(s)$, these two cases corresponding to the upper and the lower entry of (s/us) uncertainty of $\overline{m}_b$, respectively. Only in the MA/APT case, where the exact renormalon cancellation was not achieved, the parameter $k_{s/us}$ was varied between 0.1 and 10, i.e., $k_{s/us} = 1.0^{+9.0}_{-0.9}$.

---

[19] It turns out that the Landau cut in the perturbative coupling $a_{\mathrm{pt}}(\mu^2)$ starts in the 4-loop ($N_f = 3$) $\overline{\mathrm{MS}}$ scheme already at about $\mu = \Lambda_{\mathrm{L}} \approx 0.607$ GeV; while in the Lambert scheme [Eq. (10) with the central value $c_2 = -4.76$] it starts at a significantly lower scale $\mu = \Lambda_{\mathrm{L}} \approx 0.262$ GeV. The 2danQCD coupling $\mathcal{A}_1(\mu^2)$ has, of course, no Landau singularities, although it almost coincides with the Lambert pQCD coupling $a_{\mathrm{pt}}(\mu^2)$ at higher scales $|\mu^2| > 1$ GeV^2.

The other uncertainty in the determination of $\overline{m}_b$ comes from the uncertainty of the Λ^2 scale. In MA/APT it comes from $\overline{\Lambda}_5 = 0.260 \pm 0.030$ GeV and is denoted by the subscript (Λ^2) in Eq. (66). In 2danQCD and in the two pQCD approaches (Lambert scheme and $\overline{\rm MS}$ scheme), this uncertainty comes from $\alpha_s(M_Z^2; \overline{\rm MS}) = 0.1184 \pm 0.007$ [44] and is denoted by the subscript (α_s) in Eqs. (68), (70), (72).

Yet another uncertainty of $\overline{m}_b$, in 2danQCD and in Lambert scheme pQCD, comes from the variation of the (Lambert) renormalization scheme parameter $c_2 = -4.76^{+2.66}_{-0.97}$, cf. Table I.[20] and is denoted in Eqs. (68) and (70) by the subscript (c_2). On the other hand, the scheme in MA/APT was fixed by the underlying pQCD solution, Eqs. (10)-(13), with $c_2 = c_2(\overline{\rm MS}, N_f = 3) \approx +4.471$ (and $c_j = c_2^{j-1}/c_1^{j-2}$ for $j \geq 3$). In (4-loop) $\overline{\rm MS}$ pQCD, the scheme parameters are fixed in the usual way: $c_2 = c_2(\overline{\rm MS}, N_f = 3) \approx +4.471$; $c_3 = c_3(\overline{\rm MS}, N_f = 3) \approx +20.99$; $c_j = 0$ for $j \geq 4$.

When we vary the soft renormalization scale μ_s^2 around the central value $(Q_s^2)_{\rm centr.}$ of the soft reference scale [determined by Eqs. (54), (55) and (57) in pQCD, 2danQCD and MA/APT, respectively, for central values of the input parameters $\overline{m}_b$, $k_{s/us}$, etc.], so that $\mu_s^2 = 2(Q_s^2)_{\rm centr.}$ or $\mu_s^2 = (1/2)(Q_s^2)_{\rm centr.}$, this results in the corresponding variations of extracted $\overline{m}_b$ as denoted by the subscript (μ_s^2) in Eqs. (66)-(72).

Finally, the uncertainty $\delta m_b(m_c \neq 0) = \pm 0.005$ GeV due to nonzero c mass, Eq. (40) [cf. also Eqs. (46)-(49)] results in the uncertainties ∓ 0.005 GeV of $\overline{m}_b$, denoted by the subscript (m_c) in Eqs. (66)-(72).

We see in Eqs. (66)-(73) that the largest resulting uncertainty in the determination of $\overline{m}_b$ is the one originating from the uncertainty of the determination of the us binding energy (except in MA/APT where $|E_{b\bar{b}}(us)|$ values are small). These uncertainties are larger in the two pQCD approaches, due to the influence of the (unphysical) Landau singularities in the running couplings. Furthermore, the effect of the us sector in 2danQCD and in the pQCD approaches was to increase the predicted value of $\overline{m}_b$, because the us binding energies are in these cases significant and negative, cf. Fig. 3(a). If we had ignored the existence of the us sector, i.e., if we had used in the entire binding energy $E_{b\bar{b}}$ simply a common soft renormalization scale $\mu \sim Q_s$, the predicted values of $\overline{m}_b$ in 2danQCD and in Lambert and $\overline{\rm MS}$ pQCD approaches would have decreased, by -0.068, -0.091, and -0.177 GeV, respectively, as can be deduced from the us-origin uncertainties in Eqs. (68)-(72). On the other hand, in MA/APT the choice $\mu_{us} = Q_{s,({\rm MA})}$ would only slightly increase the central value of $\overline{m}_b$ by 0.002 GeV, cf. Eq. (66), basically because the values of $|E_{b\bar{b}}(us)|$ in MA/APT are much smaller, cf. Figs. 3(a),(b).

For better visibility, we present the results for the central extracted values of $\overline{m}_b$ of the aforementioned four methods in Table II, and for various uncertainties $\delta\overline{m}_b$ in Table III.

TABLE II: The extracted central values of $\overline{m}_b$ in the four models, for the central input parameter values (with the total uncertainties $\delta\overline{m}_b$). Included are the corresponding input parameter $k_{s/us}$, and the resulting scales: soft reference scale Q_s (soft renormalization scale is taken $\mu_s = Q_s$); soft mass $M_{\Upsilon(1S)}(s)$; averaged ultrasoft energy $\bar{E}_{b\bar{b}}(us)$ and its ambiguity $\delta\bar{E}_{b\bar{b}}(us)$ [cf. Eqs. (82)-(85)]. All the scales are given in GeV. Note that $M_{\Upsilon(1S)}(s) + \bar{E}_{b\bar{b}}(us) = 9.460$ GeV, i.e., the physical mass $M_{\Upsilon(1S)}$.

model	$\overline{m}_b(\delta\overline{m}_b)$	$k_{s/us}$	Q_s	$M_{\Upsilon(1S)}(s)$	$\bar{E}_{b\bar{b}}(us)$	$\delta E_{b\bar{b}}$
MA/APT	4.165(±0.022)	1.000	1.599	9.454	0.006	∓0.003
2danQCD	4.353(±0.084)	0.241	2.084	9.814	-0.353	±0.151
pQCD Lamb.	4.382(±0.111)	0.306	1.729	9.816	-0.355	±0.201
pQCD $\overline{\rm MS}$	4.505(±0.231)	0.248	1.869	10.078	-0.617	±0.394

We wish to address here briefly the question of nonperturbative (NP, higher-twist) contribution to the quarkonium mass. In the heavy quark system such as $b\bar{b}$, the nonperturbative (NP) contribution can be estimated, and in the leading order it comes from the gluon condensate and is given by [78]

$$E_{b\bar{b}}(us)^{(\rm NP)} \approx \overline{m}_b \pi^2 \frac{624}{425} \left(\frac{4\pi}{3} \overline{m}_b \right)^{-4} \frac{1}{a_{\rm pt}^4(\mu_{us}^2)} \langle \frac{\alpha_s}{\pi} G_{\mu\nu} G^{\mu\nu} \rangle \ . \tag{86}$$

---

[20] We recall that in 2danQCD all the other dimensionless parameters vary with c_2 according to Table I: the four relations $s_j(s_0)$ and $f_j(s_0)$ (for $j = 1, 2$) are fixed from the (four) conditions $\mathcal{A}_1(Q^2; c_2) - a_{\rm pt}(Q^2; c_2) \sim (\Lambda^2/Q^2)^5$ for $|Q^2| > \Lambda^2$; the scale Λ is fixed by the condition $\alpha_s(M_Z^2; \overline{\rm MS}) = 0.1184$; and the parameter s_0 $(\equiv M_0^2/\Lambda^2)$ is then fixed by the condition $r_\tau(V + A) = 0.203$. For details, we refer to Subsec. II B. Higher Lambert scheme parameters are fixed by: $c_j = c_2^{j-1}/c_1^{j-2}$ for $j \geq 3$, cf. Eqs. (10)-(13).

TABLE III: The uncertainties $\delta\overline{m}_b$ of the extracted value of $\overline{m}_b$ coming from various sources: (1) from the evaluation of the us sector; (2) from the variation of the s-us separation parameter $k_{s/us}$; (3) from the variation of α_s (or, in MA/APT: variation of Λ); (4) from the variation of the c_2 parameter (in 2danQCD, and in pQCD in the Lambert scheme); (4) from the variation of the soft renormalizatio scale μ_s; (5) from the uncertainty of $(\delta m_b)_{\rm m_c}$ of Eq. (40). See the text for details.

model	$\delta\overline{m}_b(us)$	$\delta\overline{m}_b(s/us)$	$(k_{s/us})$	$\delta\overline{m}_b(\alpha_s)$	$\delta\overline{m}_b(c_2)$	(c_2)	$\delta\overline{m}_b(\mu_s^2)$	$\delta\overline{m}_b(m_c)$
MA/	+0.002	+0.005	(1.0+9.0)	-0.019	–	(–)	-0.005	-0.005
APT	-0.002	-0.004	(1.0-0.9)	+0.020	–	(–)	+0.003	+0.005
2dan-	-0.068	+0.015	(0.241-0.100)	-0.005	-0.023	(-4.76+2.66)	+0.017	-0.005
QCD	+0.071	-0.016	(0.241+0.173)	+0.005	+0.034	(-4.76-0.97)	-0.025	+0.005
pQCD	-0.091	-0.013	(0.306+0.194)	+0.010	+0.027	(-4.76+2.66)	-0.002	-0.005
Lamb.	+0.097	+0.017	(0.306-0.119)	-0.010	-0.008	(-4.76-0.97)	-0.041	+0.005
pQCD	-0.177	-0.082	(0.248+0.642)	+0.031	–	(–)	-0.004	-0.005
$\overline{\rm MS}$	+0.200	+0.084	(0.248-0.181)	-0.027	–	(–)	-0.075	+0.005

The factor $1/a_{\rm pt}^4(\mu_{us}^2)$ in (4-loop) $\overline{\rm MS}$ pQCD is unreliable for realistic us scales $\mu_{us} \approx 1$ GeV, due to the vicinity of the Landau singularities (cf. footnote 19). In 2danQCD, for purposes of estimation, we replace $1/a_{\rm pt}^4(\mu_{us}^2)$ by $1/\mathcal{A}_4(\mu_{us}^2)$ or by $1/\widetilde{\mathcal{A}}_4(\mu_{us}^2)$. In the interval (1.1 GeV $< \mu_{us} <$ 1.3 GeV) we have $\mathcal{A}_4 \sim \widetilde{\mathcal{A}}_4 \sim 10^{-4}$; the couplings $\mathcal{A}_4$ and $\widetilde{\mathcal{A}}_4$ cover in this interval the values between 0.5×10^{-4} and 2×10^{-4}. For these values, and using the central value of the gluon condensate $\langle (\alpha_s/\pi) G^2 \rangle = 0.009$ GeV^2 [79] (cf. also Refs. [80] and [33]), and for $\overline{m}_b = 4.3$ GeV, we obtain the following estimate:

$$E_{b\bar{b}}(us)^{\rm (NP)} = 0.05^{+0.05}_{-0.02} \ {\rm GeV} \ . \tag{87}$$

This effect is relatively small and has large uncertainties. If we take it into account, then the central extracted values of $\overline{m}_b$ in this Subsection decrease somewhat, the decrease being $(\triangle \overline{m}_b)_{\rm (NP)} \approx (-0.025^{-0.025}_{+0.010})$ GeV.

We wish to dwell also on the following aspect: as seen from the results of this Subsection, the extracted values $\overline{m}_b$ and various uncertainties $\delta\overline{m}_b$ are similar in 2danQCD and the corresponding Lambert pQCD. The main reason for this is the fact that the scheme parameters (c_j, $j \geq 2$) are the same in both frameworks, and that the two corresponding running couplings practically merge at high momenta $|Q^2| > \Lambda^2$: $\mathcal{A}_1(Q^2) - a_{\rm pt}(Q^2) \sim (\Lambda^2/Q^2)^5$. On the other hand, the method of evaluation of the binding energies $E_{q\bar{q}}(s)$ and $E_{q\bar{q}}(us)$ and of the pole mass m_q was different in the two approaches. In (Lambert) pQCD, these quantities were evaluated as truncated power series in powers of $a_{\rm pt}$ [the latter determined by Eq. (10)], and the terms $a_{\rm pt}^n \ln^k a_{\rm pt}$ in $E_{q\bar{q}}$ were also evaluated directly by using the corresponding value of $a_{\rm pt}$ – cf. Eqs. (47) and (46). On the other hand, the model 2danQCD is a case of general analytic QCD, with the basic coupling $\mathcal{A}_1(Q^2)$ given in Eq. (18), and the evaluations of the higher power analogs $(a_{\rm pt}^n)_{\rm an} \equiv \mathcal{A}_n$ and of the logarithmic term analogs $(a_{\rm pt}^n \ln^k a_{\rm pt})_{\rm an} \equiv \mathcal{A}_{n,k}$ are more complicated in such a case, because linear combinations of the logarithmic derivatives $\widetilde{\mathcal{A}}_\nu$ are needed for such a construction [$\widetilde{\mathcal{A}}_\nu$ for integer ν is given in Eq. (27), and for noninteger ν in Eq. (31)] – cf. Eqs. (28)-(30) and (34). These latter relations involve truncations. For terms in $E_{q\bar{q}}$, Eqs. (50) and (61)-(64), we truncated the mentioned relations (28)-(30) and (34) at $\widetilde{\mathcal{A}}_5$ ($\sim a_{\rm pt}^5$) because perturbation expansion of $E_{q\bar{q}}$ is known up to $\sim a_{\rm pt}^5$ terms; for terms in the pole mass m_q, Eq. (53), we truncated the relations (28)-(30) at $\widetilde{\mathcal{A}}_4$ ($\sim a_{\rm pt}^4$) because perturbation expansion of m_q is known up to $\sim a_{\rm pt}^4$ terms [when the estimated r_3 term, Eq. (45), is included]. We expect that the difference between the results $\overline{m}_b$ (and $\delta\overline{m}_b$) in the 2danQCD and in Lambert scheme pQCD would decrease even more if we used in both frameworks the same method of evaluation. For example, in (Lambert) pQCD we could use for the powers of $a_{\rm pt}^n$ the linear combinations Eqs. (24)-(26) in terms of the logarithmic derivatives $\widetilde{a}_{\rm pt,n}$, Eq. (20), where the linear combinations Eqs. (24)-(26) would be truncated at $\widetilde{a}_{\rm pt,5}$ for the powers in $E_{q\bar{q}}$, and at $\widetilde{a}_{\rm pt,4}$ for powers in m_q. We note that the logarithmic derivatives in Lambert pQCD and 2danQCD differ at large momenta ($|Q^2| > \Lambda^2$) also negligibly; namely, if k is integer, we can apply $k-1$ times the derivatives $Q^2 d/(dQ^2)$ to the relation $\mathcal{A}_1(Q^2) - a_{\rm pt}(Q^2) \sim (\Lambda^2/Q^2)^5$ and obtain $\widetilde{\mathcal{A}}_k(Q^2) - \widetilde{a}_{\rm pt,k}(Q^2) \sim (\Lambda^2/Q^2)^5$ (cf. also Figs. 1-2). There is a nontrivial technical problem, though, in the evaluation of the logarithmic terms $a_{\rm pt}^n \ln^k a_{\rm pt}$ in $E_{q\bar{q}}$ along these lines in (Lambert) pQCD – for this, we would need, according to Eq. (34), to evaluate derivatives of the type $\partial^{k-s} \widetilde{a}_{{\rm pt},\nu+m}/\partial \nu^{k-s}$ at $\nu = n$ integer, but this implies that $\widetilde{a}_{{\rm pt},\nu+m}$ is needed also for ν noninteger (since variation of ν around n is needed), implying that we would need to know the logarithmic derivatives $\widetilde{a}_{{\rm pt},\nu}$ at noninteger ν, i.e., a relation of the type (31) would be needed for pQCD. This can be probably achieved by using in the integral (31) the pQCD discontinuity function for $\rho_1(\sigma)$ and extending the integration down to $\sigma = -\Lambda_{\rm L}^2 - 0$, i.e., down to the beginning of the (unphysical) Landau cut, in the spirit of Eq. (1). In the present work, we did not perform this calculation for (Lambert scheme) pQCD, though.

B. Charm mass extraction

In this case, $q\bar{q} = c\bar{c}$, the quarkonium mass is now $M_{J/\psi(1S)} = 3.0969$ GeV [44]. We basically repeat the analysis as in the case of $b\bar{b}$. There are some differences, though:

- The relation (41) is now at $N_f = 3$; therefore, the transition to the relation at the soft renormalization scales, Eq. (46), which is also at $N_f = 3$, has now no threshold transition complication.
- The typical soft reference scales [Eqs. (54) and (56)] are now significantly lower, $Q_s \approx 1$ GeV or even lower (in the $b\bar{b}$ case we had: $Q_s \approx 2$ GeV). This, in conjunction with our suggestion that the considered models 2danQCD and pQCD (in the Lambert and $\overline{\rm MS}$ schemes) are not necessarily to be trusted at scales below 1.1 GeV, implies that the typical soft renormalization scales μ_s should be chosen significantly higher than Q_s. We choose $\mu_s^2 = (3 \pm 1) Q_s^2$ in these three models. In this case, the lowest possible soft scale, $(\mu_s)_{\rm min} = \sqrt{2} Q_s$ is slightly above 1.1 GeV. In MA/APT, the soft renormalization scale will also be varied in this way, giving, however, somewhat lower central value for the renormalization scale, $\mu_s = \sqrt{3} Q_{s,({\rm MA})} \approx 1.004$ GeV.[21] In general, by keeping now the soft renormalization scale μ_s higher than the (low) soft reference scale Q_s, the uncertainties $(\delta \overline{m}_c)_{(\mu_s^2)}$ of the extracted mass $\overline{m}_c$ due to the scale variation $\mu_s^2 = (3 \pm 1) Q_s^2$ may be underestimated.
- In general, the cancellation of the leading renormalon now implies for the *s*-*us* separation parameter $k_{s/us}$ such values for which the absolute values of the *us* binding energy $E_{c\bar{c}}(us)$ are significantly smaller than in the $b\bar{b}$ case, and consequently, the *us* ambiguities are smaller. Since we now use for the central choice of the soft renormalization scale $\mu_s^2 = 3 Q_s^2$, the *us* binding energy is calculated in 2danQCD and in pQCD in the following way:

$$E_{c\bar{c}}(us) = \frac{1}{2} \left[E_{c\bar{c}}(us; 3Q_s^2) + E_{c\bar{c}}(us; 1.1^2 {\rm GeV}^2) \right] \pm \frac{1}{2} \left[E_{c\bar{c}}(us; 3Q_s^2) - E_{c\bar{c}}(us; 1.1^2 {\rm GeV}^2) \right] \ , \tag{88}$$

  where in 2danQCD the soft reference scale Q_s is determined via Eq. (56)[22] and in pQCD $Q_s = Q_{s,{\rm pt}}$ is determined via Eq. (54). In MA/APT, we do not appear to have practical problems at low scales $\mu < 1.1$ GeV, and the *us* energy as function of low scales μ_{us} turns out to have a moderate local maximum and a moderate local minimum; hence we use the formula

$$E_{c\bar{c}}(us; {\rm MA/APT}) = \frac{1}{2} \left[E_{c\bar{c}}(us)_{\rm max} + E_{c\bar{c}}(us)_{\rm min} \right] \pm \frac{1}{2} \left[E_{b\bar{b}}(us)_{\rm max} - E_{c\bar{c}}(us)_{\rm min} \right] \ , \tag{89}$$

  where these values are quite small: in the central case ($\overline{m}_c = 1.262$, $k_{s/us} = 1.0$), the local maximum is $E_{c\bar{c}}(us)_{\rm max} \approx +0.003$ GeV and is reached at $\mu_{us} \approx 0.69$ GeV, and the local minimum is $E_{c\bar{c}}(us)_{\rm min} \approx -0.005$ GeV and is reached at very low scale $\mu_{us} \approx 0.10$ GeV.
- The exact renormalon cancellation requirement in MA/APT now does not give an exceedingly small value of the *s*-*us* separation parameter $k_{s/us}$, but it it still quite small, $k_{s/us} \approx 0.02$. We decide, for presentation reasons, to vary in MA/APT the parameter $k_{s/us}$ again around its central chosen value 1.0, in the interval between 0.1 and 10., just as it was done in the $b\bar{b}$ case of MA/APT.

The described behavior of the $E_{c\bar{c}}(us; \mu_{us}^2)$ in the analytic 2danQCD model and in pQCD in the two schemes, as a function of the ulttrasoft scales μ_{us}, is presented in Fig. 4(a), and in the case of MA/APT in Fig. 4(b). Comparing Figs. 4(a) and (b), we see that $|E_{c\bar{c}}(us)| \sim 10^{-2}$-$10^{-1}$ GeV in 2danQCD, and $\sim 10^{-3}$ GeV in MA/APT. Furthermore, comparing with the corresponding curves in Figs. 3(a) and (b) for the $b\bar{b}$, we can see that in 2danQCD and in the two pQCD approaches, the absolute values $|E_{c\bar{c}}(us)|$ are by almost two orders of magnitude smaller than $|E_{b\bar{b}}(us)|$, principally because the renormalon cancellation gives us in the two cases significantly different *s*-*us* separation parameter values: $k_{s/us}(c\bar{c}) = 4.00$ and $k_{s/us}(b\bar{b}) = 0.241$. Further, in general the values of $|E_{q\bar{q}}(us)|$ in pQCD are larger than in 2danQCD, this being true even more so in $\overline{\rm MS}$ pQCD than in Lambert scheme pQCD, basically because of the relatively high branch cut scale $\Lambda_{\rm L}^2$ at which the Landau singularities cut start in the $\overline{\rm MS}$ case (cf. footnote 19). In the MA/APT case, the values of $|E_{q\bar{q}}(us)|$ are quite small in both cases ($q = c, b$), being in $c\bar{c}$ case smaller by almost one order of magnitude. All this is reflected also in the numerical results of this and of the previous Subsection.

---

[21] If we used in MA/APT a lower definition of the central renormalization scale, $\mu_s = Q_{s,({\rm MA})}$ (≈ 0.58 GeV), the predicted central value of $\overline{m}_c$ would go up by only 0.005 GeV.

[22] Note that in 2danQCD in the $c\bar{c}$ case the condition (55) cannot be fulfilled.



FIG. 4: (a) The ultrasoft binding energy $E_{c\bar{c}}(us)$ (in GeV) as a function of the us renormalization scale μ_{us}, for the central input values of 2danQCD model for the $J/\psi(1S)$ system ($\overline{m}_c = 1,266$ GeV; $k_{s/us} = 4.00$; $c_2 = -4.76$; $s_0 = 23.06$): solid line for 2danQCD; dashed line for Lambert pQCD ($c_2 = -4.76$); dash-dotted line for the 4-loop $\overline{\mathrm{MS}}$ pQCD ($c_2 = 4.471$). For all three curves the same $\overline{m}_c$ and $k_{s/us}$ values are used. Included are the properly rescaled 2danQCD couplings $\mathcal{A}_5(\mu_{us}^2)$ and $\mathcal{A}_{5,1}(\mu_{us}^2)$ as functions of the scale μ_{us}. (b) The same as in Figure (a), but for the MA/APT model (with $c_2 = 4.471$), with the corresponding central values of that model ($\overline{m}_c = 1.262$ GeV and $k_{s/us} = 1$). In all the curves $N_f = 3$ was taken.

The resulting extracted values of $\overline{m}_c$ are

$$\begin{aligned}\overline{m}_c(\mathrm{MA/APT}) &= \left\{1.262 \mp 0.002_{(us)} \mp 0.002_{(s/us)} \mp 0.011_{(\Lambda^2)} \mp 0.002_{(\mu_s^2)}\right\} \mathrm{GeV} &\quad (90)\\ &= 1.262 \pm 0.012 \ \mathrm{GeV}\ , \quad \text{with}: \sqrt{3}Q_{s,(\mathrm{MA})} = 1.00 \ \mathrm{GeV}\ ,\ k_{s/us} = 1.0\ . &\quad (91)\end{aligned}$$

$$\begin{aligned}\overline{m}_c(\mathrm{2danQCD}) &= \left\{1.266 \pm 0.003_{(us)} \pm 0.007_{(s/us)} \mp 0.005_{(\alpha_s)} + \begin{pmatrix} -0.014 \\ +0.003 \end{pmatrix}_{(c_2)} + \begin{pmatrix} -0.002 \\ +0.005 \end{pmatrix}_{(\mu_s^2)} \right\} \mathrm{GeV} &\quad (92)\\ &= 1.266 \pm 0.017 \ \mathrm{GeV}\ , \quad \text{with}: \sqrt{3}Q_s = 1.42 \ \mathrm{GeV}\ ,\ k_{s/us} = 4.0\ . &\quad (93)\end{aligned}$$

$$\begin{aligned}\overline{m}_c(\mathrm{pQCDLamb.}) &= \left\{1.265 \pm 0.001_{(us)} + \begin{pmatrix} +0.021 \\ -0.021 \end{pmatrix}_{(s/us)} \mp 0.004_{(\alpha_s)} \pm 0.000_{(c_2)} + \begin{pmatrix} -0.003 \\ +0.015 \end{pmatrix}_{(\mu_s^2)} \right\} \mathrm{GeV} &\quad (94)\\ &= 1.265 \pm 0.027 \ \mathrm{GeV}\ , \quad \text{with}: \sqrt{3}Q_s = 1.38 \ \mathrm{GeV}\ ,\ k_{s/us} = 5.59\ . &\quad (95)\end{aligned}$$

$$\begin{aligned}\overline{m}_c(\mathrm{pQCD}\overline{\mathrm{MS}}) &= \left\{1.272 \mp 0.011_{(us)} + \begin{pmatrix} +0.066 \\ -0.075 \end{pmatrix}_{(s/us)} \mp 0.002_{(\alpha_s)} + \begin{pmatrix} -0.003 \\ +0.017 \end{pmatrix}_{(\mu_s^2)} \right\} \mathrm{GeV} &\quad (96)\\ &= 1.272 \pm 0.078 \ \mathrm{GeV}\ , \quad \text{with}: \sqrt{3}Q_s = 1.58 \ \mathrm{GeV}\ ,\ k_{s/us} = 3.08\ . &\quad (97)\end{aligned}$$

In order to see more clearly the renormalon cancellation, we present below, as in Subsec. IV A, the sum for the soft mass $M_{J/\psi(1S)}(s)$, combining in each parenthesis the (positive) term $\sim a^n$ from $2m_c(\overline{m}_c^2; \mu_s^2)$ and the (negative) term $\sim a^{n+1}$ from the soft binding energy $E_{c\bar{c}}(s; \overline{m}_c^2; \mu_s^2)$ ($n = 0, 1, \ldots, 4$), for the central input values of parameters $\overline{m}_c$, μ_s ($= \sqrt{3}Q_s$) and $k_{s/us}$; separately we present below also $E_{c\bar{c}}(us)$.

$$\begin{aligned}M_{J/\psi(1S)}(s; \mathrm{MA/APT}) &= 2.524 + (0.412 - 0.056) + (0.227 - 0.081) + (0.125 - 0.071) + (0.045 - 0.026) \ \mathrm{GeV} & \\ &= 2.524 + 0.356 + 0.146 + 0.054 + 0.020 \ \mathrm{GeV}\ (= 3.099 \ \mathrm{GeV})\ , &\quad (98)\\ E_{c\bar{c}}(us; \mathrm{MA/APT}) &= -0.001 \pm 0.004 \ \mathrm{GeV}\ , \qquad (\overline{m}_c = 1.262 \ \mathrm{GeV},\ k_{s/us} = 1.0)\ ; &\quad (99)\end{aligned}$$

$$\begin{aligned}M_{J/\psi(1S)}(s; \mathrm{2danQCD}) &= 2.532 + (0.349 - 0.053) + (0.328 - 0.171) + (0.253 - 0.150) + (0.979 - 0.979) \ \mathrm{GeV} & \\ &= 2.532 + 0.296 + 0.157 + 0.103 + 0.000 \ \mathrm{GeV}\ (= 3.088 \ \mathrm{GeV})\ , &\quad (100)\\ E_{c\bar{c}}(us; \mathrm{2danQCD}) &= 0.009 \mp 0.006 \ \mathrm{GeV}\ , \qquad (\overline{m}_c = 1.266 \ \mathrm{GeV},\ k_{s/us} = 4.0)\ ; &\quad (101)\end{aligned}$$

$$\begin{aligned}M_{J/\psi(1S)}(s; \mathrm{pQCDLamb.}) &= 2.530 + (0.354 - 0.061) + (0.303 - 0.141) + (0.404 - 0.305) + (0.662 - 0.662) \ \mathrm{GeV} & \\ &= 2.530 + 0.293 + 0.161 + 0.099 + 0.000 \ \mathrm{GeV}\ (= 3.083 \ \mathrm{GeV})\ , &\quad (102)\\ E_{c\bar{c}}(us; \mathrm{pQCDLamb.}) &= 0.013 \mp 0.003 \ \mathrm{GeV}\ , \qquad (\overline{m}_c = 1.265 \ \mathrm{GeV},\ k_{s/us} = 5.59)\ ; &\quad (103)\end{aligned}$$

TABLE IV: The extracted central values of $\overline{m}_c$ in the four models, for the central input parameter values (with the total uncertainties $\delta\overline{m}_c$). Included are the corresponding input parameter $k_{s/us}$, and the resulting scales: soft renormalization scale $\mu_s = \sqrt{3}Q_s$; soft mass $M_{J/\psi(1S)}(s)$; averaged ultasoft energy $\bar{E}_{c\bar{c}}(us)$ and its ambiguity $\delta\bar{E}_{c\bar{c}}(us)$ [cf. Eqs. (88)-(89)]. All the scales are given in GeV. Note that $M_{J/\psi(1S)}(s) + \bar{E}_{c\bar{c}}(us) = 3.097$ GeV, i.e., the physical mass $M_{J/\psi(1S)}$.

model	$\overline{m}_c(\delta\overline{m}_c)$	$k_{s/us}$	$\sqrt{3}Q_s$	$M_{J/\psi(1S)}(s)$	$\bar{E}_{c\bar{c}}(us)$	$\delta E_{c\bar{c}}$
MA/APT	1.262(±0.012)	1.00	1.004	3.099	-0.001	±0.004
2danQCD	1.266(±0.017)	4.00	1.423	3.088	+0.009	∓0.006
pQCD Lamb.	1.265(±0.027)	5.59	1.382	3.083	+0.013	∓0.003
pQCD $\overline{\rm MS}$	1.272(±0.078)	3.08	1.585	3.131	-0.034	±0.024

TABLE V: The uncertainties $\delta\overline{m}_c$ of the extracted value of $\overline{m}_c$ coming from various sources: (1) from the evaluation of the us sector; (2) from the variation of the s-us separation parameter $k_{s/us}$; (3) from the variation of α_s (or, in MA/APT: variation of Λ); (4) from the variation of the c_2 parameter (in 2danQCD, and in pQCD in the Lambert scheme); (4) from the variation of the soft renormalizatio scale μ_s; See the text for details.

model	$\delta\overline{m}_c(us)$	$\delta\overline{m}_c(s/us)$	$(k_{s/us})$	$\delta\overline{m}_c(\alpha_s)$	$\delta\overline{m}_c(c_2)$	(c_2)	$\delta\overline{m}_c(\mu_s^2)$
MA/	-0.002	-0.002	(1.0+9.0)	-0.011	–	(–)	-0.002
APT	+0.002	+0.002	(1.0-0.9)	+0.011	–	(–)	+0.002
2dan-	+0.003	+0.007	(4.00-3.61)	-0.005	-0.014	(-4.76+2.66)	-0.002
QCD	-0.003	-0.007	(4.00+39.5)	+0.005	+0.003	(-4.76-0.97)	+0.005
pQCD	+0.001	+0.021	(5.59-5.355)	-0.004	0.000	(-4.76+2.66)	-0.003
Lamb.	-0.001	-0.022	(5.59+148.4)	+0.004	0.000	(-4.76-0.97)	+0.015
pQCD	-0.011	+0.066	(3.08-2.86)	-0.002	–	(–)	-0.003
$\overline{\rm MS}$	+0.011	-0.075	(3.08+66.9)	+0.002	–	(–)	+0.017

$$\begin{aligned}
M_{J/\psi(1S)}(s; {\rm pQCD}\overline{\rm MS}) &= 2.544 + (0.368 - 0.066) + (0.348 - 0.163) + (0.455 - 0.356) + (0.778 - 0.778) \text{ GeV} \\
&= 2.544 + 0.302 + 0.185 + 0.099 + 0.000 \text{ GeV } (= 3.131 \text{ GeV}) \ , \qquad (104) \\
E_{c\bar{c}}(us; {\rm pQCD}\overline{\rm MS}) &= -0.034 \pm 0.024 \text{ GeV} \ , \qquad (\overline{m}_c = 1.272 \text{ GeV}, \ k_{s/us} = 3.08) \ ; \qquad (105)
\end{aligned}$$

As in the previous Subsection in the case of $b\bar{b}$, we present now for the case $c\bar{c}$, for better visibility, the results for the central extracted values of $\overline{m}_c$ of the four models in Table IV, and for various uncertainties $\delta\overline{m}_c$ in Table V.

The nonperturbative (NP) contribution coming from the gluon condensate, cf. Eq. (86) for the $b\bar{b}$ system in the previous Subsection, is for the lighter $c\bar{c}$ system unreliable, since the next-to-leading corrections are in this case large and tend to make the result unreliable, cf. Ref. [81].

V. SUMMARY

We evaluated, in two analytic QCD models and in perturbative QCD (pQCD, in two schemes), the quark-antiquark binding energies (up to N^{3}LO) and masses of ground state $q\bar{q}$ quarkonia ($q = b, c$), as functions of the quark mass $\overline{m}_q$ ($\equiv \overline{m}_q(\mu^2 = \overline{m}_q^2)$), also called $\overline{\rm MS}$ quark mass. In analytic QCD models the QCD running coupling $\mathcal{A}_1(Q^2)$ has no unphysical (Landau) singularities in the Q^2-plane. One analytic model used was the Minimal Analytic model (referred to also as: Analytic Perturbation Theory, MA/APT) of Shirkov, Solovtsov, Solovtsova, and Milton, Refs. [11–14]. The (analytic) QCD running coupling in this model, $\mathcal{A}_1^{\rm (MA)}(Q^2)$, differs nonnegligibly from the corresponding (same scheme) pQCD coupling $a_{\rm pt}(Q^2) \equiv \alpha_s(Q^2)/\pi$ even at large momenta $|Q^2| > \Lambda^2$: $\mathcal{A}_1^{\rm (MA)}(Q^2) - a_{\rm pt}(Q^2) \sim (\Lambda^2/Q^2)^1$. The other analytic QCD model used was the so called 2-delta analytic QCD (2danQCD), Ref. [32], which basically coincides with pQCD in the same (Lambert) scheme at high momenta $|Q^2| > \Lambda^2$: $\mathcal{A}_1(Q^2) - a_{\rm pt}(Q^2) \sim (\Lambda^2/Q^2)^5$. The use of the analytic QCD models was motivated by the fact that the typical soft (s) momentum scales Q_s in the ground bound states of quarkonia are low ($Q_s \approx 2$ GeV and 1 GeV, for $b\bar{b}$ and $c\bar{c}$, respectively), and that the typical ultrasoft (us) momentum scales Q_{us} are even lower. This, in conjunction with the fact that Landau singularities of the pQCD coupling $a_{\rm pt}(Q^2)$ reach relatively high momenta: $Q \approx 0.61$ GeV in the usual (4-loop) $\overline{\rm MS}$ scheme (with $c_2 \equiv \beta_2/\beta_0 = 4.471$), and $Q \approx 0.26$ GeV in the Lambert scheme ($c_2 = -4.76$).

In the course of the calculations, we included a renormalon-based estimate of the N^{3}LO term ($\sim a_{\rm pt}^4$) of the quark pole mass m_q in terms of $a_{\rm pt}$ and $\overline{m}_q$, cf. Appendix B. This allowed us to fix the s-us separation of the binding energy, $E_{q\bar{q}} = E_{q\bar{q}}(s) + E_{q\bar{q}}(us)$, via the requirement of cancellation of the leading infrared renormalon in the sum $M_{q\bar{q}}(s) = 2m_q + E_{q\bar{q}}(s)$. In the latter sum we used as the input parameter the mass $\overline{m}_q$ which is free of the renormalon ambiguity, and the coefficients of the perturbation expansion in $a_{\rm pt}(\mu_s^2)$ at the same common soft renormalization scale μ_s^2, where for the quark pole mass and for the soft binding energy terms up to $\sim a_{\rm pt}^4$ and $\sim a_{\rm pt}^5$ were used, respectively. It turned out that in MA/APT the renormalon cancellation condition cannot be exactly, but only approximately, fulfilled. In the case of the $b\bar{b}$ system ($\Upsilon(1s)$), this condition gave in 2danQCD and in pQCD the values of the *s*-*us* separation parameter $k_{s/us}$, Eq. (59), between 0.24 and 0.31. This led to significant (and negative) values of the *us* part $E_{b\bar{b}}(us)$ of the binding energy, cf. Fig. 3(a). In pQCD these values had a particularly large uncertainties due to the vicinity of Landau singularities. These effects of the *us* regime led, in comparison with the usual perturbative approaches in the literature, to higher central values of the bottom quark mass: $\overline{m}_b \approx 4.35, 4.38, 4.51$ GeV in 2danQCD, Lambert pQCD, $\overline{\rm MS}$ pQCD, respectively. Further, these effects led to large uncertainties, especially in $\overline{\rm MS}$ pQCD – cf. Tables II and III. If we used in $E_{b\bar{b}}(us)$ the same (soft) renormalization scale as used in $E_{b\bar{b}}(s)$ ($\mu \sim Q_s \sim \alpha_s \overline{m}_b$), the extracted central values of $\overline{m}_b$ would decrease, by about −0.07, −0.09, −0.18 GeV, respectively. The 2danQCD analytic model gave us, among other things, clues about how to estimate the large negative values of $E_{b\bar{b}}(us)$ and the uncertainties thereof, not just in 2danQCD, but also in pQCD in both schemes. The other, less significant, uncertainties in the extraction of the value of $\overline{m}_b$, come from the uncertainty of the *s*-*us* separation parameter $k_{s/us}$, from the uncertainty of the world average of the QCD coupling $\alpha_s(M_Z^2; \overline{\rm MS})$, from the renormalization scheme and renormalization (soft) scale variations (c_2 and μ_s^2), and from the uncertainty of the finite charm quark mass effect at μ_s scales. On the other hand, MA/APT has in general quite suppressed couplings in the infrared, so that the *us* sector does not affect it significantly. Our results for $\overline{m}_b$ are given in Eqs. (66)-(73), and in Tables II and III.

In the $c\bar{c}$ system ($J/\psi(1S)$), the analysis was similar. Surprisingly, the infrared renormalon cancellation condition gave in this case for the *s*-*us* separation parameter $k_{s/us}$ significantly higher values, between 3.1 and 5.6 in 2danQCD and in pQCD (in MA/APT the cancellation condition was again not achieved). As a result, despite having in general even lower typical *us* scales μ_{us} than in the $b\bar{b}$ case, the *us* parts $E_{c\bar{c}}(us)$ of the binding energies have in general significantly smaller absolute values than in the $b\bar{b}$ case, cf. Figs. 4(a) and 3(a). This resulted in smaller influence of the *us* sector on the extraction of $\overline{m}_c$, and also in smaller uncertainties $\delta\overline{m}_c$ from this sector, than in the case of $b\bar{b}$ ($\overline{m}_b$ and $\delta\overline{m}_b$) – cf. Eqs. (90)-(97), and Tables IV and V. The obtained central extracted values of $\overline{m}_c \approx 1.27$ GeV are thus close to those in the literature. In MA/APT the results for $\overline{m}_c$ are also similar.

The main conclusion of this work appears to be the following: analytic QCD approaches which at high energies follow the pQCD behavior closely (such as 2danQCD analytic QCD) indicate that the ultrasoft regime in the $b\bar{b}$ system is important. Our approach indicates that the influence of the ultrasoft regime is underestimated in the usual pQCD approaches (especially in the $\overline{\rm MS}$ scheme) which consider the soft and ultrasoft energies to be roughly equal and consequently give lower values of the extracted masses $\overline{m}_b$ ($\overline{m}_b < 4.3$ GeV).

Acknowledgments

We thank A. Pineda for useful comments. A.C. thanks J. Otálora for help in using calculational software. This work was supported in part by Mecesup2 (Chile) Grant FSM 0605-D3021 (C.A.), and by FONDECYT (Chile) Grant No. 1095196 and Anillos Project ACT119 (G.C.).

Appendix A: Ground state quark-antiquark binding energy

According to Refs. [7, 58, 64], the perturbation expansion of the quark-antiquark binding energy $E_{q\bar{q}}$, in terms of the quark pole mass m_q and $\alpha_s(\mu^2) \equiv \alpha_s$, is

$$E_{q\bar{q}} = E_1^C + \delta E_1^{(1)} + \delta E_1^{(2)} + \delta E_1^{(3)} \ , \tag{A1}$$

where

$$E_1^C \ = \ -\frac{1}{4}\alpha_s^2 C_F^2 m \ ; \tag{A2}$$

$$\delta E_1^{(1)} \ = \ -\frac{\alpha_s^3 \beta_0 C_F^2 m}{\pi} \left[-\ln\left(\frac{\alpha_s C_F m}{\mu}\right) + \psi(2) \right] - \frac{2\alpha_s^3 C_F^2 m}{\pi} \left(a_1 + \frac{\beta_0 \gamma_E}{2} \right) \ ; \tag{A3}$$

$$\begin{aligned}
\delta E_1^{(2)} &= -\frac{4\alpha_s^4\beta_0^2C_F^2m}{\pi^2}\left[N_0-N_1\ln\left(\frac{\alpha_sC_Fm}{\mu}\right)+\frac{1}{4}\ln^2\left(\frac{\alpha_sC_Fm}{\mu}\right)\right]+\delta_{s1}\frac{\alpha_s^4C_F^4m}{3}+\frac{a_2\alpha_s^4C_F^3m}{4}\\
&-\frac{2\alpha_s^4\beta_0^2C_F^2m}{\pi^2}\left[\ln^2\left(\frac{\alpha_sC_Fm}{\mu}\right)-2\ln\left(\frac{\alpha_sC_Fm}{\mu}\right)\psi(2)+\psi(2)+\psi(2)^2+\psi'(2)\right.\\
&\left.+\frac{2\Gamma(1)}{\Gamma(2)}\theta(-1)\sum_{j=1}^{-1}\frac{\Gamma(2+j)}{j!j^2}\right]-\frac{\alpha_s^4c_2^{(L)}C_F^2m}{2\pi^2}\left[-\ln\left(\frac{\alpha_sC_Fm}{\mu}\right)+\psi(2)\right]\\
&-\frac{4\alpha_s^4\beta_0C_F^2m}{\pi^2}\left[-\left(a_1+\frac{\beta_0\gamma_E}{2}\right)+\left(a_1+\frac{\beta_0\gamma_E}{2}\right)\left(-\ln[\frac{\alpha_sC_Fm}{\mu}]+\psi(2)\right)\right]\\
&-\frac{5\alpha_s^4C_F^4m}{8^2}-\frac{3\alpha_s^4C_F^4m}{8}-\frac{4\alpha_s^4\beta_0C_F^2m}{\pi^2}\left[\left(a_1+\frac{\beta_0\gamma_E}{2}\right)^2\right.\\
&\left.+2\left(b_1+\left(a_1\beta_0+\frac{\beta_1}{8}\right)\gamma_E+\frac{\beta_0^2}{4}\left(\gamma_E^2+\frac{\pi^2}{12}\right)\right)\right]\ ; \qquad \text{(A4)}
\end{aligned}$$

$$\delta E_1^{(3)} = \delta E_1^{(3)}|_{\beta(\alpha_s)=0}+\delta E_1^{(3)}|_{\beta(\alpha_s)}\ . \qquad \text{(A5)}$$

The two contributions to $\delta E_1^{(3)}$ are

$$\begin{aligned}
\delta E_1^{(3)}|_{\beta(\alpha_s)=0} &= -E_1^C\frac{\alpha_s^3}{\pi}\left\{-\frac{a_1\tilde{a}_2+a_3}{32\pi^2}+a_1\left[-\frac{C_AC_F}{2}+C_F^2\left(-\frac{19}{16}+\frac{S(1+S)}{2}\right)\right]\right.\\
&+C_A^3\left(-\frac{1}{36}+\frac{L_{\alpha_s}}{6}+\frac{\ln[2]}{6}\right)+C_A^2C_F\left(-\frac{49}{36}+\frac{4}{3}\left(L_{\alpha_s}+\ln[2]\right)\right)\\
&+C_AC_F^2\left[-\frac{5}{72}+\frac{37L_{\alpha_s}}{6}+\left(\frac{85}{54}-\frac{7}{6}L_{\alpha_s}\right)S(1+S)+\frac{10\ln[2]}{3}\right]\\
&+C_F^3\left(\frac{50}{9}+3L_{\alpha_s}-\frac{S(1+S)}{3}+\frac{8\ln[2]}{3}\right)+C_F^2T_F\left(-\frac{32}{15}+S(1+S)(1-\ln[2])\right.\\
&\left.\left.+2\ln[2]\right)+\frac{49}{36}C_AC_FN_fT_F+C_F^2N_fT_F\left(\frac{11}{18}-\frac{10}{27}S(1+S)\right)+\frac{2}{3}C_F^3L_1^E\right\}\ , \qquad \text{(A6)}
\end{aligned}$$

$$\begin{aligned}
\delta E_1^{(3)}|_{\beta(\alpha_s)} &= E_1^C\frac{\alpha_s^3}{\pi^3}\left\{32\beta_0^3L_\mu^3+\left[12a_1\beta_0^2+40\beta_0^3+28\beta_0\beta_1\right]L_\mu^2\right.\\
&+L_\mu\left[10a_1\beta_0^2+3a_1\beta_1+4\beta_2+\beta_0\left(\frac{a_1^2}{2}+\tilde{a}_2+40\beta_1+8C_AC_F\pi^2\right.\right.\\
&\left.\left.+C_F^2\left(\frac{21\pi^2}{2}-\frac{16}{3}\pi^2S(1+S)\right)\right)+\beta_0^3\left(\frac{16\pi^2}{3}+64\zeta(3)\right)\right]+\beta_0^3\left(-8+4\pi^2+\frac{2\pi^4}{45}\right.\\
&\left.+64\zeta(3)-8\pi^2\zeta(3)+96\zeta(5)\right)+a_1\beta_0^2\left(\frac{2\pi^2}{3}+8\zeta(3)\right)+\beta_0\left[-\frac{a_1^2}{8}+\frac{3\tilde{a}_2}{4}\right.\\
&+C_AC_F\left(6\pi^2-\frac{2\pi^4}{3}\right)+C_F^2\left(8\pi^2-\frac{4\pi^4}{3}+\left(-\frac{4\pi^2}{3}+\frac{4\pi^4}{9}\right)S(1+S)\right)\\
&\left.\left.+\beta_1\left(8+\frac{7\pi^2}{3}+16\zeta(3)\right)\right]+2a_1\beta_1+4\beta_2\right\}\ . \qquad \text{(A7)}
\end{aligned}$$

The following notations were used:

$$\begin{aligned}
L_\mu &= \ln\left[\frac{\mu}{\alpha_sC_Fm}\right]\ , \qquad L_{\alpha_s}=-\ln\left[C_F\alpha_s\right]\ , \qquad L_1^E=-81.5379\ ,\\
&S=1\ , \qquad c_2^{(L)}=a_1\beta_0+\frac{\beta_1}{8}+\frac{\beta_0^2\gamma_E}{2}\ , \qquad a_2=\frac{1}{2}\left(-2C_A+C_F\right)\ ,\\
N_0 &= \frac{1}{12}\left(3+3{\gamma_E}^2-\pi^2+6\zeta(3)\right)\ , \qquad N_1=-\frac{\gamma_E}{2}\ , \qquad \text{(A8)}\\
C_A &= 3\ , \qquad C_F=4/3\ , \qquad N_f=1,2,3\ , \qquad T_F=1/2\ . \qquad \text{(A9)}
\end{aligned}$$

The RGE coefficients β_j are in $\overline{\rm MS}$ scheme

$$\begin{aligned}
\beta_0 &= \frac{1}{4}\left(11 - \frac{2}{3}N_f\right) \ , \qquad \beta_1 = \frac{1}{16}\left(102 - \frac{38}{3}N_f\right) \ , \\
\beta_2 &= \frac{1}{64}\left(\frac{2857}{2} - \frac{5033}{18}N_f + \frac{325}{54}{N_f}^2\right) \ , \\
\beta_3 &= \frac{1}{256}\left[\frac{149753}{6} + \frac{1093}{729}{N_f}^3 + 3564\zeta(3) + {N_f}^2\left(\frac{50065}{162} + \frac{6472\zeta(3)}{81}\right)\right. \\
& \left. -N_f\left(\frac{1078361}{162} + \frac{6508\zeta(3)}{27}\right)\right] \ .
\end{aligned} \tag{A10}$$

The constants a_1 and $\tilde{a}_2$ are

$$\begin{aligned}
a_1 &= \frac{1}{9}\left(31C_A - 20N_f T_F\right) \ , \\
\tilde{a}_2 &= \frac{400{N_f}^2{T_F}^2}{81} - C_F N_f T_F\left(\frac{55}{3} - 16\zeta(3)\right) + C_A^2\left(\frac{4343}{162} + \frac{1}{4}\left(16\pi^2 - \pi^4\right) + \frac{22\zeta(3)}{3}\right) \\
& -C_A N_f T_F\left(\frac{1798}{81} + \frac{56\zeta(3)}{3}\right) \ .
\end{aligned} \tag{A11}$$

The value for the constant a_3 asociated with the three-loop soft contribution was obtained in Refs. [5, 6] and is

$$a_3 = a_3^{(0)} + a_3^{(1)}N_f + a_3^{(2)}N_f^2 + a_3^{(3)}N_f^3 \ , \tag{A12}$$

where

$$\begin{aligned}
a_3^{(0)} &= 502.24C_A^3 - 136.39\left(\frac{N_C\left(N_C^2+6\right)}{48}\right) \ , \\
a_3^{(1)} &= -709.717C_A^2T_F + \left(\frac{-71281}{162} + 264\zeta(3) + 80\zeta(5)\right)C_AC_FT_F \\
& + \left(\frac{286}{9} + 296\frac{\zeta(3)}{3} - 160\zeta(5)\right)C_F^2T_F - 56.83\left(\frac{18 - 6N_C^2 + N_C^4}{96N_C^2}\right) \ , \\
a_3^{(2)} &= C_FT_F^2\left(\frac{14002}{81} - \frac{416\zeta(3)}{3}\right) + C_AT_F^2\left(\frac{12541}{243} + \frac{64\pi^4}{135} + \frac{368\zeta(3)}{3}\right) \ , \\
a_3^{(3)} &= -\frac{8000T_F^3}{729} \ .
\end{aligned} \tag{A13}$$

Appendix B: Renormalon-based estimate of $\sim a_{\rm pt}^4$ coefficient

The term r_3 in the expansion of $m_q/\overline{m}_q$ in Eq. (41) can be estimated by a method closely related with the approach presented in Sec. II of Ref. [74]. The pQCD version of the sum in Eq. (41) can be reexpressed in terms of $a_{\rm pt}(\mu^2)$ at any other renormalization scale μ^2

$$S \equiv \frac{m_q}{\overline{m}_q} - 1 = \frac{4}{3}a_{\rm pt}(\mu^2)\left[1 + a_{\rm pt}(\mu^2)r_1(\mu^2) + a_{\rm pt}^2(\mu^2)r_2(\mu^2) + \mathcal{O}(a_{\rm pt}^3)\right] \ , \tag{B1a}$$

$$r_1(\mu^2) = \kappa_1 + \beta_0 L_m(\mu^2) \ , \tag{B1b}$$

$$r_2(\mu^2) = \kappa_2 + (2\kappa_1\beta_0 + \beta_1)L_m(\mu^2) + \beta_0^2 L_m^2(\mu^2) \ , \tag{B1c}$$

$$(4/3)\kappa_1 = 6.248\beta_0 - 3.739 \ , \tag{B1d}$$

$$(4/3)\kappa_2 = 23.497\beta_0^2 + 6.248\beta_1 + 1.019\beta_0 - 29.94 \ , \tag{B1e}$$

where $L_m(\mu^2) = \ln(\mu^2/\overline{m}_q^2)$, while $\beta_0(N_f)$ and $\beta_1(N_f)$ are the renormalization scheme independent coefficients (given just after Eq. (13)). Here, $N_f = N_\ell$ is the number of light active flavors (quarks with masses lighter than m_q).

Since r_1 and r_2 are explicitly known, the Borel transform $B_S(b)$ is known to order $\sim b^2$

$$B_S(b;\mu) = \frac{4}{3}\left[1 + \frac{r_1(\mu^2)}{1!\,\beta_0} b + \frac{r_2(\mu^2)}{2!\,\beta_0^2} b^2 + \mathcal{O}(b^3)\right] \ . \tag{B2}$$

The function $B_S(b)$ has renormalon singularities at $b = 1/2, 3/2, 2, \ldots, -1, -2, \ldots$ [82–84]. The behavior of B_S near the leading infrared (IR) renormalon singularity $b = 1/2$ is determined by the resulting renormalon ambiguity of m_q. This ambiguity δm_q is a (QCD) scale which, having the dimension of energy and being renormalization scale and scheme indepenent, must be proportional to the QCD scale $\Lambda_{\rm QCD}$: $\delta m_q = const \times \Lambda_{\rm QCD}$ [85]. This scale can be expressed in terms of $a_{\rm pt}(\mu^2)$ and μ (μ being any renormalization scale) in the form

$$\Lambda = const \times \mu \exp\left(-\frac{1}{2\beta_0 a_{\rm pt}(\mu)}\right) a_{\rm pt}(\mu)^{-\nu} c_1^{-\nu} \left[1 + \sum_{k=1}^{\infty} (2\beta_0)^k \nu(\nu-1)\cdots(\nu-k+1)\widetilde{c}_k a_{\rm pt}^k(\mu)\right] \ , \tag{B3}$$

where

$$\nu \ = \ \frac{c_1}{2\beta_0} = \frac{\beta_1}{2\beta_0^2} \ , \tag{B4a}$$

$$\widetilde{c}_1 \ = \ \frac{(c_1^2 - c_2)}{(2\beta_0)^2 \nu} \ , \qquad \widetilde{c}_2 = \frac{1}{2(2\beta_0)^4 \nu(\nu-1)} \left[(c_1^2 - c_2)^2 - 2\beta_0(c_1^3 - 2c_1c_2 + c_3)\right] \ , \tag{B4b}$$

$$\widetilde{c}_3 \ = \ \frac{1}{6(2\beta_0)^6 \nu(\nu-1)(\nu-2)} \left[(c_1^2 - c_2)^3 - 6\beta_0(c_1^2 - c_2)(c_1^3 - 2c_1c_2 + c_3) + 8\beta_0^2(c_1^4 - 3c_1^2c_2 + c_2^2 + 2c_1c_3 - c_4)\right] \tag{B4c}$$

The above constants, expressed in terms of β_0 and of $c_j = \beta_j/\beta_0$, appear in the expansion of the residue of the Borel transform $B_S(b)$ at the pole $b = 1/2$

$$B_S(b;\mu) \ = \ N_m \pi \frac{\mu}{\overline{m}_q} \frac{1}{(1-2b)^{1+\nu}} \left[1 + \sum_{k=1}^{\infty} \widetilde{c}_k (1-2b)^k\right] + B_S^{(\rm an.)}(b;\mu) \ , \tag{B5}$$

where $B_S^{(\rm an.)}(b;\mu)$ is analytic on the disk $|b| < 1$ and can be expanded in powers of b We can assume that the coefficients $\widetilde{c}_k$ are known up to $k = 3$, because the coefficient $c_4 = \beta_4/\beta_0$ (in $\overline{\rm MS}$ scheme) is known to a large degree by Padé-related methods of Ref. [86]

$$\beta_4 = \frac{1}{4^5}(A_4 + B_4 N_f + C_4 N_f^2 + D_4 N_f^3 + E_4 N_f^4) \tag{B6}$$

with $A_4 = 7.59 \times 10^5$, $B_4 = -2.19 \times 10^5$, $C_4 = 2.05 \times 10^4$, $D_4 = -49.8$, and $E_4 = -1.84$. This gives $c_4 = 123.7$ for $N_f = 3$, $c_4 = 97.2$ for $N_f = 4$, and $c_4 = 86.2$ for $N_f = 5$. The residue parameter N_m can be determined with high precision by using the idea of Refs. [87], i.e., by calculating (cf. Refs. [72, 73, 88]):

$$N_m = \frac{\overline{m}_q}{\mu} \frac{1}{\pi} R_S(b = 1/2) \ , \tag{B7}$$

where

$$R_S(b;\mu) \equiv (1-2b)^{1+\nu} B_S(b;\mu) \ , \tag{B8}$$

and the first coefficients in the expansion in powers of b of this quantity are known from the known coefficients r_1 and r_2. We can use a combination of truncated perturbation series and Padé approximants [1/1] for $R_S(b)$, as presented in Ref. [88], and obtain

$$N_m \ \approx \ 0.575 (N_f = 3) \ , \quad \approx 0.555 (N_f = 4) \ , \quad \approx 0.533 (N_f = 5) \ . \tag{B9}$$

with the uncertainties in these values of roughly ±0.020.

In the bilocal expansion (B5), the analytic part $B_S^{(\rm an.)}(b;\mu)$ can be taken as a polynomial in b, i.e., a truncated expansion in powers of b. The coefficients of the latter expansion can be related with $r_j(\mu^2)$'s by equating the expansion of Eq. (B5) in powers of b with the expansion (B2), resulting in

$$B_S^{(\rm an.)}(b;\mu) \ = \ h_0^{(m)} + \sum_{k \geq 1} \frac{h_k^{(m)}}{k!\,\beta_0^k} b^k \ , \tag{B10a}$$

$$h_k^{(m)} \ = \ \frac{4}{3} r_k - \pi N_m \frac{\mu}{\overline{m}_q} (2\beta_0)^k \sum_{n \geq 0} \widetilde{c}_n \frac{\Gamma(\nu + k + 1 - n)}{\Gamma(\nu + 1 - n)} \ , \tag{B10b}$$

where, by convention, $r_0 = \widetilde{c}_0 = 1$. The numbers $\widetilde{c}_n$ of Eqs. (B4), which enter the sum in Eq. (B10b), are known only up to $n = 3$, because, in $\overline{\rm MS}$, only c_k up to $k = 4$ are reasonably known (c_4 approximately, as mentioned). For $N_f = 3$, these values are: $\widetilde{c}_1 = -0.1638$, $\widetilde{c}_2 = 0.2372$, $\widetilde{c}_3 = -0.1205$ (and: $\nu = 0.3951$). For $N_f = 4$, they are: $\widetilde{c}_1 = -0.1054$, $\widetilde{c}_2 = 0.2736$, $\widetilde{c}_3 = -0.1610$ (and: $\nu = 0.3696$). And for $N_f = 5$ they are: $\widetilde{c}_1 = 0.0238$, $\widetilde{c}_2 = 0.3265$, $\widetilde{c}_3 = -0.2681$ (and: $\nu = 0.3289$). Therefore, the sums in (B10b) are truncated at $n = 3$.

Theoretically, the pole closest to the origin in $B_S^{\rm (an.)}(b;\mu)$ is at $b = -1$, at least in the large-β_0 approximation.[23] Since in $B_S^{\rm (an.)}(b;\mu)$ the coefficients $h_k^{(m)}(\mu^2)$ for $k = 0, 1, 2$ are known (because $r_1(\mu^2)$ and $r_2(\mu^2)$ are known), we can construct the Padé $[1/1]_{B_S^{\rm (an.)}}(b)$ and check the pole of it. It turns out that this Padé, at the natural scale $\mu = \overline{m}_q$, has the pole at $b = -1.09, -0.96, -1.12$, for $N_f = 3, 4, 5$, respectively, reflecting correctly the theoretical expectation.[24] We can extend further this reasoning and obtain the next coefficient $h_3^{(m)}$ (at $\mu = \overline{m}_q$) by requiring that the Padé $[2/1]_{B_S^{\rm (an.)}}(b)$ have the pole at $b = -1$. This gives us

$$h_3^{(m)}(\overline{m}_q^2) = -25.18 (N_f = 3) \ , \ -28.28 (N_f = 4) \ , \ -35.62 (N_f = 5) \ . \tag{B11}$$

If constructing with these values of $h_3^{(m)}$ the other possible Padé approximant of index 3, namely $[1/2]_{B_S^{\rm (an.)}}(b)$, it turns out that the nearest to origin pole of such Padé is then at $b = -1.003, -1.001, -1.008$, for $N_f = 3, 4, 5$, respectively. This indicates that the obtained values of $h_3^{(m)}$, Eq. (B11), are consistent.[25] Using these values, we obtain from the relations (B10b) (with the natural choice $\mu^2 = \overline{m}_q^2$) the estimate for r_3

$$r_3(\overline{m}_q^2) \ = \ \frac{3}{4} h_k^{(m)}(\overline{m}_q^2) + \frac{3\pi}{4} N_m (2\beta_0)^k \sum_{n=0}^{3} \widetilde{c}_n \frac{\Gamma(\nu + k + 1 - n)}{\Gamma(\nu + 1 - n)} \ . \tag{B12}$$

This gives us the values for $r_3 \equiv r_3(\overline{m}_q^2)$ the following estimates:

$$r_3 = 1339.4 (N_f = 3) \ , \ 987.3 (N_f = 4) \ , \ 690.1 (N_f = 5) \ . \tag{B13}$$

The principal origin of the uncertainties in these expressions is the uncertainty in the residue parameter N_m (roughly ± 0.020, i.e., less than 4 %), implying an uncertainty in r_3 of a few percent (below 4 %).

An analysis similar to this one has been performed in Ref. [72]. There, however, the term $\widetilde{c}_3$ and the coefficients $h_k^{(m)}$ were not included in the analysis.

---

[23] Nonetheless, there is a possibility that at two-loop order the kinetic term contributes to an IR renormalon at $b = +1$ in $B_S(b)$, cf. Ref. [91].

[24] Though, the μ-dependence of this position is rather strong. E.g., when μ^2 varies by 10 % around $\overline{m}_q^2$, the pole position in [1/1] varies between -1.6 and -0.7 in $N_f = 3$ case, between -1.2 and -0.7 in $N_f = 4$ case, and between -1.26 and -1.00 in $N_f = 5$ case.

[25] We used for $\overline{\rm MS}$ scheme coefficient c_4 the estimated values (B6), with $c_4 = 123.7, 97.2, 86.2$, for $N_f = 3, 4, 5$, respectively, from Ref. [86]. Simpler Padé-based estimates of c_4 were obtained in Ref. [89]: $c_4 = 40, 70$ for $N_f = 4, 5$, respectively (and a large negative and uncertain value $c_4 = -850$ for $N_f = 3$). The $c_4 = 40$ value (for $N_f = 4$) in this case differs substantially from the value $c_4 = 97.2$. If we repeat for the $c_4 = 40$ value ($N_f = 4$) the same procedure described above, we obtain $\widetilde{c}_4 \approx 0.0055$ (for $c_4 = 97.2$ we got: $\widetilde{c}_4 = -0.1610$), hence the expressions of $h_k^{(m)}(\overline{m}_q^2)$ of Eq. (B10b) change, and the pole of $[1/1]_{B_S^{\rm (an.)}}(b)$ becomes $b \approx -5.9$ (before: $b \approx -0.96$), not close to the theoretical pole $b = -1$. Furthermore, from the requirement that $[2/1]_{B_S^{\rm (an.)}}(b)$ have the pole at $b = -1$ we now get $h_3^{(m)}(\overline{m}_q^2) = 4.10$ (before: -28.28), and using this value of $h_3^{(m)}$ in the Padé $[1/2]_{B_S^{\rm (an.)}}(b)$ we obtain the pole nearest to the origin $b = 1.94$ (before: $b = -1.001$). This indicates that the estimate $c_4 = 40$ (for $N_f = 4$) is not giving results consistent with the theoretical expectations of the renormalon structure of $B_S^{\rm (an.)}$.

Appendix C: Variation of pQCD coupling with scales and schemes

In this Appendix we give the relation between $a_0 \equiv a_{\rm pt}(Q_0^2; c_2^{(0)}, c_3^{(0)}, \ldots)$ and $a \equiv a_{\rm pt}(Q^2; c_2, c_3, \ldots)$, where the latter is expressed as power expansion of the former (cf. Appendix A of Ref. [90] for details)

$$\begin{aligned} a &= a_0 + a_0^2(-x) + a_0^3(x^2 - c_1 x + \delta c_2) \\ &\quad + a_0^4(-x^3 + \frac{5}{2} c_1 x^2 - c_2^{(0)} x - 3x\delta c_2 + \frac{1}{2}\delta c_3) \\ &\quad + a_0^5 \Big[x^4 - \frac{13}{3} c_1 x^3 + (\frac{3}{2} c_1^2 + 3 c_2^{(0)} + 6\delta c_2) x^2 \\ &\quad + (-c_3^{(0)} - 3 c_1 \delta c_2 - 2\delta c_3) x + (\frac{1}{3} c_2^{(0)} \delta c_2 + \frac{5}{3}(\delta c_2)^2 - \frac{1}{6} c_1 \delta c_3 + \frac{1}{3}\delta c_4) \Big] + \mathcal{O}(a_0^6) \ , \end{aligned} \tag{C1}$$

where we denote

$$a \equiv a_{\rm pt}(Q^2; c_2, c_3, \ldots) \ , \quad a_0 \equiv a_{\rm pt}(Q_0^2; c_2^{(0)}, c_3^{(0)}, \ldots) \ , \tag{C2}$$

$$x \equiv \beta_0 \ln \frac{Q^2}{Q_0^2}, \quad \delta c_k \equiv c_k - c_k^{(0)} \ . \tag{C3}$$

For the purposes of our paper, it is sufficient to consider in the above relation (C1) terms up to (including) terms $\sim a_0^4$.

The 3-loop threshold connection of $a_{\rm pt}$ in $\overline{\rm MS}$ scheme at the threshold scale $\mu_{\rm thr}^2 = (\mathcal{K} \overline{m}_c)^2$ (where $\mathcal{K} \sim 1$; usually $\mathcal{K} = 2$) can be written as the following relation between $a_{\rm pt}(\mu_{\rm thr}^2 + 0; N_f = 4) \equiv a_+$ and $a_{\rm pt}(\mu_{\rm thr}^2 - 0; N_f = 3) \equiv a_-$:

$$a_+ = a_- \left[1 + x_1 a_- + x_2 a_-^2 + x_3 a_-^3 + \mathcal{O}(a_-^4) \right] \ , \tag{C4}$$

where

$$x_1 = -k_1 \ , \qquad x_2 = -k_2 + 2k_1^2 \ , \qquad x_3 = -k_3 + 5k_1 k_2 - 5k_1^3 \ , \tag{C5}$$

and the coefficients k_j were calculated in Ref. [66]

$$\begin{aligned} k_1 &= -\frac{1}{6}\ell_h \ , \qquad k_2 = \frac{1}{36}\ell_h^2 - \frac{19}{24}\ell_h + \frac{11}{72} \ , \\ k_3 &= -\frac{1}{216}\ell_h^3 - \frac{131}{576}\ell_h^2 + \frac{(-6793 + 281 N_\ell)}{1728}\ell_h + \left(-\frac{82043}{27648}\zeta(3) + \frac{564731}{124416} - \frac{2633}{31104} N_\ell \right) \ , \end{aligned} \tag{C6}$$

where $\ell_h = \ln(\mu_{\rm thr}^2/\overline{m}_c^2) = \ln \mathcal{K}^2$, and N_ℓ in Eq. (C6) is the number of light quark flavors, i.e., $N_\ell = 3$ in the considered case of transition from $N_f = 4$ to $N_f = 3$.

[1] M. Peter, Phys. Rev. Lett. **78** (1997) 602 [arXiv:hep-ph/9610209]; Nucl. Phys. B **501** (1997) 471 [arXiv:hep-ph/9702245].

[2] Y. Schröder, Phys. Lett. B **447** (1999) 321 [arXiv:hep-ph/9812205]; Nucl. Phys. Proc. Suppl. **86** (2000) 525 [arXiv:hep-ph/9909520].

[3] N. Brambilla, A. Pineda, J. Soto and A. Vairo, Phys. Rev. D **60** (1999) 091502 [arXiv:hep-ph/9903355];

[4] B. A. Kniehl and A. A. Penin, Nucl. Phys. B **563** (1999) 200 [arXiv:hep-ph/9907489].

[5] A. V. Smirnov, V. A. Smirnov and M. Steinhauser, Phys. Rev. Lett. **104**, 112002 (2010) [arXiv:0911.4742 [hep-ph]].

[6] C. Anzai, Y. Kiyo and Y. Sumino, Phys. Rev. Lett. **104**, 112003 (2010) [arXiv:0911.4335 [hep-ph]].

[7] A. A. Penin and M. Steinhauser, Phys. Lett. B **538** (2002) 335 [arXiv:hep-ph/0204290].

[8] A. Pineda, Prog. Part. Nucl. Phys. **67**, 735 (2012) [arXiv:1111.0165 [hep-ph]].

[9] N.N. Bogoliubov and D.V. Shirkov, *Introduction to the Theory of Quantum Fields*, New York, Wiley, 1959; 1980.

[10] R. Oehme, Int. J. Mod. Phys. A **10**, 1995 (1995) [arXiv:hep-th/9412040].

[11] D. V. Shirkov and I. L. Solovtsov, hep-ph/9604363; Phys. Rev. Lett. **79**, 1209 (1997) [arXiv:hep-ph/9704333].

[12] K. A. Milton, I. L. Solovtsov and O. P. Solovtsova, Phys. Lett. B **415**, 104 (1997) [arXiv:hep-ph/9706409].

[13] D. V. Shirkov, Theor. Math. Phys. **127**, 409 (2001) [hep-ph/0012283];

[14] D. V. Shirkov, Eur. Phys. J. C **22**, 331 (2001) [hep-ph/0107282].

[15] A. V. Nesterenko, Phys. Rev. D **62**, 094028 (2000); Phys. Rev. D **64**, 116009 (2001); Int. J. Mod. Phys. A **18**, 5475 (2003);

[16] A. V. Nesterenko and J. Papavassiliou, Phys. Rev. D **71**, 016009 (2005); A. C. Aguilar, A. V. Nesterenko and J. Papavassiliou, J. Phys. G **31**, 997 (2005). J. Phys. G **32**, 1025 (2006) [arXiv:hep-ph/0511215]; A. V. Nesterenko, arXiv:0710.5878 [hep-ph].

[17] A. I. Alekseev, Few Body Syst. **40**, 57 (2006) [arXiv:hep-ph/0503242].

[18] Y. Srivastava, S. Pacetti, G. Pancheri and A. Widom, *In the Proceedings of* e^+e^- *Physics at Intermediate Energies, SLAC, Stanford, CA, USA, 30 April - 2 May 2001, pp T19* [arXiv:hep-ph/0106005].

[19] B. R. Webber, JHEP **9810**, 012 (1998) [arXiv:hep-ph/9805484].

[20] G. Cvetič and C. Valenzuela, J. Phys. G **32**, L27 (2006) [arXiv:hep-ph/0601050].

[21] G. Cvetič and C. Valenzuela, Phys. Rev. D **74**, 114030 (2006) [arXiv:hep-ph/0608256].

[22] U. Aglietti, G. Ferrera and G. Ricciardi, Nucl. Phys. B **768**, 85 (2007) [hep-ph/0608047]; U. Aglietti, F. Di Lodovico, G. Ferrera and G. Ricciardi, Eur. Phys. J. C **59**, 831 (2009) [arXiv:0711.0860 [hep-ph]].

[23] D. V. Shirkov, arXiv:1208.2103 [hep-th].

[24] G. M. Prosperi, M. Raciti and C. Simolo, Prog. Part. Nucl. Phys. **58**, 387 (2007) [arXiv:hep-ph/0607209].

[25] D. V. Shirkov and I. L. Solovtsov, Theor. Math. Phys. **150**, 132 (2007) [arXiv:hep-ph/0611229].

[26] G. Cvetič and C. Valenzuela, Braz. J. Phys. **38**, 371 (2008) [arXiv:0804.0872 [hep-ph]].

[27] A. P. Bakulev, Phys. Part. Nucl. **40**, 715 (2009) [arXiv:0805.0829 [hep-ph]] (arXiv preprint in Russian).

[28] Y. .O. Belyakova and A. V. Nesterenko, Int. J. Mod. Phys. A **26**, 981 (2011) [arXiv:1011.1148 [hep-ph]].

[29] G. Cvetič, R. Kögerler and C. Valenzuela, J. Phys. G **37**, 075001 (2010) [arXiv:0912.2466 [hep-ph]];

[30] G. Cvetič, R. Kögerler and C. Valenzuela, Phys. Rev. D **82**, 114004 (2010) [arXiv:1006.4199 [hep-ph]].

[31] C. Contreras, G. Cvetič, O. Espinosa and H. E. Martínez, Phys. Rev. D **82**, 074005 (2010) [arXiv:1006.5050].

[32] C. Ayala, C. Contreras and G. Cvetič, Phys. Rev. D **85**, 114043 (2012) [arXiv:1203.6897 [hep-ph]].

[33] G. Cvetič and C. Villavicencio, arXiv:1209.2953 [hep-ph].

[34] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147**, 385 (1979); Nucl. Phys. B **147**, 448 (1979).

[35] Y. L. Dokshitzer, G. Marchesini and B. R. Webber, Nucl. Phys. B **469**, 93 (1996) [arXiv:hep-ph/9512336].

[36] I. L. Solovtsov and D. V. Shirkov, Phys. Lett. B **442**, 344 (1998) [hep-ph/9711251].

[37] A. P. Bakulev, S. V. Mikhailov and N. G. Stefanis, Phys. Rev. D **72**, 074014 (2005) [Erratum-ibid. D **72**, 119908 (2005)] [arXiv:hep-ph/0506311].

[38] A. P. Bakulev, S. V. Mikhailov and N. G. Stefanis, Phys. Rev. D **75**, 056005 (2007) [Erratum-ibid. D **77**, 079901 (2008)] [arXiv:hep-ph/0607040].

[39] A. P. Bakulev, S. V. Mikhailov and N. G. Stefanis, JHEP **1006**, 085 (2010) [arXiv:1004.4125 [hep-ph]].

[40] D. V. Shirkov, Nucl. Phys. Proc. Suppl. **162**, 33 (2006) [arXiv:hep-ph/0611048].

[41] G. Cvetič and A. V. Kotikov, J. Phys. G **39**, 065005 (2012) [arXiv:1106.4275 [hep-ph]].

[42] P. A. Baikov, K. G. Chetyrkin and J. H. Kühn, Phys. Rev. Lett. **101**, 012002 (2008) [arXiv:0801.1821 [hep-ph]].

[43] A. P. Bakulev, private communication.

[44] K. Nakamura *et al.* [Particle Data Group Collaboration], J. Phys. G **37**, 075021 (2010); J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86**, 010001 (2012).

[45] E. Gardi, G. Grunberg and M. Karliner, JHEP **9807**, 007 (1998) [hep-ph/9806462];

[46] D. S. Kourashev, arXiv:hep-ph/9912410; D. S. Kurashev and B. A. Magradze, Theor. Math. Phys. **135**, 531 (2003) [Teor. Mat. Fiz. **135**, 95 (2003)].

[47] B. A. Magradze, Few Body Syst. **40**, 71 (2006) [hep-ph/0512374].

[48] Mathematica 8.0.4, Wolfram Co.

[49] R. S. Pasechnik, D. V. Shirkov and O. V. Teryaev, Phys. Rev. D **78**, 071902 (2008) [arXiv:0808.0066 [hep-ph]]; V. L. Khandramai, R. S. Pasechnik, D. V. Shirkov, O. P. Solovtsova and O. V. Teryaev, Phys. Rev. D **81**, 016010 (2010) [arXiv:0911.3297]; Phys. Lett. B **706**, 340 (2012) [arXiv:1106.6352 [hep-ph]].

[50] S. Peris, M. Perrottet and E. de Rafael, JHEP **9805**, 011 (1998) [arXiv:hep-ph/9805442].

[51] B. A. Magradze, Few Body Syst. **48**, 143 (2010) [Erratum-ibid. **53**, 365 (2012)] [arXiv:1005.2674 [hep-ph]].

[52] K. A. Milton, I. L. Solovtsov and O. P. Solovtsova, Phys. Rev. D **64**, 016005 (2001) [arXiv:hep-ph/0102254].

[53] M. Baldicchi, A. V. Nesterenko, G. M. Prosperi, D. V. Shirkov and C. Simolo, Phys. Rev. Lett. **99**, 242001 (2007); M. Baldicchi, A. V. Nesterenko, G. M. Prosperi and C. Simolo, Phys. Rev. D **77**, 034013 (2008).

[54] A. V. Nesterenko, arXiv:1209.0164 [hep-ph].

[55] R. Barate *et al.* [ALEPH Collaboration], Eur. Phys. J. C **4**, 409 (1998); S. Schael *et al.* [ALEPH Collaboration], Phys. Rept. **421**, 191 (2005) [hep-ex/0506072]; M. Davier, A. Höcker and Z. Zhang, Rev. Mod. Phys. **78**, 1043 (2006) [hep-ph/0507078].

[56] M. Davier, S. Descotes-Genon, A. Höcker, B. Malaescu and Z. Zhang, Eur. Phys. J. C **56**, 305 (2008) [arXiv:0803.0979 [hep-ph]].

[57] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).

[58] A. Pineda and F. J. Yndurain, Phys. Rev. D **61** (2000) 077505; Phys. Rev. D **58** (1998) 094022.

[59] N. Brambilla, A. Pineda, J. Soto and A. Vairo, Phys. Lett. B **470**, 215 (1999) [hep-ph/9910238].

[60] N. Brambilla, Y. Sumino and A. Vairo, Phys. Rev. D **65** (2002) 034001 [arXiv:hep-ph/0108084].

[61] R. Tarrach, Nucl. Phys. B **183**, 384 (1981).

[62] N. Gray, D. J. Broadhurst, W. Grafe and K. Schilcher, Z. Phys. C **48** (1990) 673; L. V. Avdeev and M. Y. Kalmykov, Nucl. Phys. B **502**, 419 (1997) [arXiv:hep-ph/9701308].

[63] K. G. Chetyrkin and M. Steinhauser, Phys. Rev. Lett. **83**, 4001 (1999).

[64] K. G. Chetyrkin and M. Steinhauser, Nucl. Phys. B **573**, 617 (2000).

[65] K. Melnikov and T. v. Ritbergen, Phys. Lett. B **482**, 99 (2000).

[66] K. G. Chetyrkin, B. A. Kniehl and M. Steinhauser, Phys. Rev. Lett. **79**, 2184 (1997) [arXiv:hep-ph/9706430].

[67] A. H. Hoang, M. C. Smith, T. Stelzer and S. Willenbrock, Phys. Rev. D **59** (1999) 114014 [arXiv:hep-ph/9804227].

[68] N. Brambilla, A. Pineda, J. Soto and A. Vairo, Nucl. Phys. B **566** (2000) 275 [arXiv:hep-ph/9907240].

[69] M. Beneke, Phys. Lett. B **434** (1998) 115 [arXiv:hep-ph/9804241].
[70] Y. Kiyo and Y. Sumino, Phys. Lett. B **496**, 83 (2000) [hep-ph/0007251]; Phys. Rev. D **67**, 071501 (R) (2003) [arXiv:hep-ph/0211299].
[71] N. Brambilla, Y. Sumino and A. Vairo, Phys. Lett. B **513**, 381 (2001) [hep-ph/0101305].
[72] A. Pineda, JHEP **0106**, 022 (2001) [arXiv:hep-ph/0105008].
[73] T. Lee, JHEP **0310**, 044 (2003) [arXiv:hep-ph/0304185].
[74] C. Contreras, G. Cvetič and P. Gaete, Phys. Rev. D **70** (2004) 034008 [arXiv:hep-ph/0311202].
[75] A. H. Hoang, Z. Ligeti and A. V. Manohar, Phys. Rev. Lett. **82**, 277 (1999) [arXiv:hep-ph/9809423]; Phys. Rev. D **59**, 074017 (1999) [arXiv:hep-ph/9811239].
[76] D. Eiras and J. Soto, Phys. Lett. B **491**, 101 (2000) [arXiv:hep-ph/0005066].
[77] A. H. Hoang, arXiv:hep-ph/0008102.
[78] M. B. Voloshin, Nucl. Phys. B **154**, 365 (1979); H. Leutwyler, Phys. Lett. B **98**, 447 (1981); M. B. Voloshin, Sov. J. Nucl. Phys. **36**, 143 (1982) [Yad. Fiz. **36**, 247 (1982)].
[79] B. L. Ioffe and K. N. Zyablyuk, Eur. Phys. J. C **27**, 229 (2003) [arXiv:hep-ph/0207183].
[80] B. V. Geshkenbein, B. L. Ioffe and K. N. Zyablyuk, Phys. Rev. D **64**, 093009 (2001) [arXiv:hep-ph/0104048]; B. L. Ioffe, Prog. Part. Nucl. Phys. **56**, 232 (2006) [arXiv:hep-ph/0502148].
[81] A. Pineda, Nucl. Phys. B **494**, 213 (1997) [hep-ph/9611388].
[82] I. I. Bigi, M. A. Shifman, N. G. Uraltsev and A. I. Vainshtein, Phys. Rev. D **50** (1994) 2234 [arXiv:hep-ph/9402360].
[83] M. Beneke and V. M. Braun, Nucl. Phys. B **426** (1994) 301 [arXiv:hep-ph/9402364];
[84] M. Beneke, Phys. Rept. **317** (1999) 1.
[85] M. Beneke, Phys. Lett. B **344**, 341 (1995) [arXiv:hep-ph/9408380].
[86] J. R. Ellis, I. Jack, D. R. Jones, M. Karliner and M. A. Samuel, Phys. Rev. D **57**, 2665 (1998) [arXiv:hep-ph/9710302].
[87] T. Lee, Phys. Rev. D **56** (1997) 1091 [arXiv:hep-th/9611010]; Phys. Lett. B **462** (1999) 1 [arXiv:hep-ph/9908225].
[88] G. Cvetič, J. Phys. G G **30**, 863 (2004) [hep-ph/0309262].
[89] V. Elias, T. G. Steele, F. Chishtie, R. Migneron and K. B. Sprague, Phys. Rev. D **58**, 116007 (1998) [arXiv:hep-ph/9806324].
[90] G. Cvetič and R. Kögerler, Phys. Rev. D **63**, 056013 (2001) [hep-ph/0006098].
[91] M. Neubert, Phys. Lett. B **393**, 110 (1997) [hep-ph/9610471].